

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number
WO 01/09120 A1

(51) International Patent Classification⁷: C07D 401/12,
401/14, 213/40, 471/10, C07C 311/18, C07D 401/04,
211/34, 295/14, 213/38, 409/14, 333/20, A61K 31/444,
31/4523, A61P 25/00

(21) International Application Number: PCT/US00/20482

(22) International Filing Date: 27 July 2000 (27.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/146,069 28 July 1999 (28.07.1999) US

(71) Applicant: ORTHO-MCNEIL PHARMACEUTICAL,
INC. [US/US]; U.S. Route #202, P.O. Box 300, Raritan, NJ
08869-0602 (US).

(72) Inventors: DAX, Scott, L.; 3 Quail Drive, Landenberg, PA
19350 (US). MCNALLY, James; 321 Heatherfield Drive,
Souderton, PA 18964 (US). YOUNGMAN, Mark; 675 E.
Street Road, Warminster, PA 18974 (US).

(74) Agents: CIAMPORCERO, Audley, A., Jr. et al.;
Johnson & Johnson, One Johnson & Johnson Plaza, New
Brunswick, NJ 08933 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.


(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- With international search report.
- Before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: AMINE AND AMIDE DERIVATIVES AS LIGANDS FOR THE NEUROPEPTIDE Y Y5 RECEPTOR USEFUL IN
THE TREATMENT OF OBESITY AND OTHER DISORDERS



(A)

(57) Abstract: Amine and amide derivatives of formula (A) which are ligands for the neuropeptide Y Y5 (NPY5) receptor, methods of preparation and pharmaceutical compositions containing amines and amides of formula (A) as the active ingredient are described. The amines and amides of formula (A) are useful in the treatment of disorders and diseases associated with NPY receptor subtype Y5.

WO 01/09120 A1

**AMINE AND AMIDE DERIVATIVES AS LIGANDS
FOR THE NEUROPEPTIDE Y Y5 RECEPTOR
USEFUL IN THE TREATMENT OF OBESITY AND OTHER DISORDERS**

5 FIELD OF THE INVENTION

This invention relates to a series of amine and amide derivatives, pharmaceutical compositions containing them and intermediates used in their preparation. The compounds of the invention are ligands for the neuropeptide
10 Y Y5 (NPY5) receptor, a receptor which is associated with a number of central nervous system disorders and affective conditions. In addition, many of the compounds of the invention reduce food consumption in a rodent model of feeding.

15 BACKGROUND OF THE INVENTION

Regulation and function of the mammalian central nervous system is governed by a series of interdependent receptors, neurons, neurotransmitters, and proteins. The neurons play a vital role in this system, for when externally
20 or internally stimulated, they react by releasing neurotransmitters that bind to specific proteins. Common examples of endogenous small molecule neurotransmitters such as acetylcholine, adrenaline, norepinephrine, dopamine, serotonin, glutamate, and gamma-aminobutyric acid are well known, as are the specific receptors that recognize these compounds as
25 ligands ("The Biochemical Basis of Neuropharmacology", Sixth Edition, Cooper, J. R.; Bloom, F. E.; Roth, R. H. Eds., Oxford University Press, New York, NY 1991).

In addition to the endogenous small molecule neurotransmitters, there is
30 increasing evidence that neuropeptides play an integral role in neuronal operations. Neuropeptides are now believed to be co-localized with perhaps more than one-half of the 100 billion neurons of the human central nervous system. In addition to humans, neuropeptides have been discovered in a

number of animal species. In some instances the composition of these peptides is remarkably homogenous among species. This finding suggests that the function of neuropeptides is vital and has been impervious to evolutionary changes. Furthermore, neuropeptides, unlike small molecule neurotransmitters, are typically synthesized by the neuronal ribosome. In some cases, the active neuropeptides are produced as part of a larger protein which is enzymatically processed to yield the active substance. Based upon these differences, compared to small molecule neurotransmitters, neuropeptide-based strategies may offer novel therapies for CNS diseases and disorders. Specifically, agents that affect the binding of neuropeptides to their respective receptors or ameliorate responses that are mediated by neuropeptides are potential therapies for diseases associated with neuropeptides.

There are a number of afflictions that are associated with the complex interdependent system of receptors and ligands within the central nervous system; these include neurodegenerative diseases, affective disorders such as anxiety, depression, pain and schizophrenia, and affective conditions that include a metabolic component, namely obesity. Such conditions, disorders and diseases have been treated with small molecules and peptides which modulate neuronal responses to endogenous neurotransmitters.

One example of the class of neuropeptides is neuropeptide Y (NPY). NPY was first isolated from porcine brain (Tatemoto, K. et al. *Nature* 1982, 296, 659) and was shown to be structurally similar to other members of the pancreatic polypeptide (PP) family such as peptide YY, which is primarily synthesized by endocrine cells in the gut, and pancreatic polypeptide, which is synthesized by the pancreas. Neuropeptide Y is a single peptide protein that consists of thirty-six amino acids containing an amidated C-terminus. Like other members of the pancreatic polypeptide family, NPY has a distinctive conformation that consists of an N-terminal polyproline helical region and an amphiphilic α -helix joined by a characteristic PP-fold (Vladimir, S. et al. *Biochemistry* 1990, 20, 4509). Furthermore, NPY sequences from a number of

animal species have been elucidated and all show a high degree of amino acid homology to the human protein (>94% in rat, dog, rabbit, pig, cow, sheep) (see Larhammar, D. in "The Biology of Neuropeptide Y and Related Peptides", Colmers, W. F. and Wahlestedt, C. Eds., Humana Press, Totowa, NJ 1993).

5

Endogenous receptor proteins that bind NPY and related peptides as ligands have been identified and distinguished, and several such proteins have been cloned and expressed. Six different receptor subtypes [Y1, Y2, Y3, Y4(PP), Y5, Y6 (formerly designated as a Y5 receptor)] are recognized today
10 based upon binding profile, pharmacology and / or composition if identity is known (Wahlestedt, C. et. al. *Ann. NY Acad. Sci.* **1990**, 611, 7; Larhammar, D. et. al. *J. Biol. Chem.* **1992**, 267, 10935; Wahlestedt, C. et. al. *Regul. Pept.* **1986**, 13, 307; Fuhlendorff, J. U. et. al. *Proc. Natl. Acad. Sci. USA* **1990**, 87, 182; Grundemar, L. et. al. *J. Pharmacol. Exp. Ther.* **1991**, 258, 633; Laburthe,
15 M. et. al. *Endocrinology* **1986**, 118, 1910; Castan, I. et. al. *Endocrinology* **1992**, 131, 1970; Gerald, C. et. al. *Nature* **1996**, 382, 168; Weinberg, D. H. et. al. *Journal of Biological Chemistry* **1996**, 271, 16435; Gehlert, D. et. al. *Current Pharmaceutical Design* **1995**, 1, 295; Lundberg, J. M. et. al. *Trends in Pharmaceutical Sciences* **1996**, 17, 301). Most and perhaps all NPY receptor
20 proteins belong to the family of so-called G-protein coupled receptors (GPCRs). The neuropeptide Y5 receptor, a putative GPCR, is negatively coupled to cellular cyclic adenosine monophosphate (cAMP) levels via the action of adenylate cyclase (Gerald, C. et. al. *Nature* **1996**, 382, 168; Gerald, C. et. al. PCT WO 96/16542). For example, NPY inhibits forskolin-stimulated
25 cAMP production / levels in a neuroblastoma cell line. A Y5 ligand that mimics NPY in this fashion is an agonist whereas one that competitively reverses the NPY inhibition of forskolin-stimulated cAMP production is an antagonist.

Neuropeptide Y itself is the archetypal substrate for the NPY receptors
30 and its binding can elicit a variety of pharmacological and biological effects *in vitro* and *in vivo*. When administered to the brain of live animals (intracerebroventricularly (icv) or into the amygdala), NPY produces anxiolytic effects in established animal models of anxiety such as the elevated plus-

5 maze, Vogel punished drinking and Geller-Seifter's bar-pressing conflict paradigms (Heilig, M. et. al. *Psychopharmacology* **1989**, 98, 524; Heilig, M. et. al. *Reg. Peptides* **1992**, 41, 61; Heilig, M. et. al. *Neuropsychopharmacology* **1993**, 8, 357). Thus compounds that mimic NPY are postulated to be useful for the treatment of anxiolytic disorders.

10 The immunoreactivity of neuropeptide Y is notably decreased in the cerebrospinal fluid of patients with major depression and those of suicide victims (Widdowson, P. S. et. al. *Journal of Neurochemistry* **1992**, 59, 73), and rats treated with tricyclic antidepressants display significant increases of NPY relative to a control group (Heilig, M. et. al. *European Journal of Pharmacology* **1988**, 147, 465). These findings suggest that an inadequate NPY response may play a role in some depressive illnesses, and that compounds that regulate the NPY-ergic system may be useful for the treatment of depression.

15

Neuropeptide Y improves memory and performance scores in animal models of learning (Flood, J. F. et. al. *Brain Research* **1987**, 421, 280) and therefore may serve as a cognition enhancer for the treatment of neurodegenerative diseases such as Alzheimer's Disease (AD) as well as
20 AIDS-related and senile dementia.

Elevated plasma levels of NPY are present in animals and humans experiencing episodes of high sympathetic nerve activity such as surgery, newborn delivery and hemorrhage (Morris, M. J. et. al. *Journal of Autonomic Nervous System* **1986**, 17, 143). Thus chemical substances that alter the
25 NPY-ergic system may be useful for alleviating migraine, pain and the condition of stress.

Neuropeptide Y also mediates endocrine functions such as the release
30 of luteinizing hormone (LH) in rodents (Kalra, S. P. et. al. *Frontiers in Neuroendocrinology* **1992**, 13, 1). Since LH is vital for mammalian ovulation, a compound that mimics the action of NPY could be useful for the treatment of infertility, particularly in women with so-called luteal phase defects.

Neuropeptide Y is a powerful stimulant of food intake; as little as one-billionth of a gram, when injected directly into the CNS, causes satiated rats to overeat (Clark, J. T. et. al. *Endocrinology* **1984**, 115, 427; Levine, A. S. et. al. *Peptides* **1984**, 5, 1025; Stanley, B. G. et. al. *Life Sci.* **1984**, 35, 2635; Stanley, B. G. et. al. *Proc. Nat. Acad. Sci. USA* **1985**, 82, 3940). Thus NPY is orexigenic in rodents but not anxiogenic when given intracerebroventricularly and so antagonism of neuropeptide receptors may be useful for the treatment of diabetes and eating disorders such as obesity, anorexia nervosa and bulimia nervosa.

In recent years, a variety of potent, structurally distinct small molecule Y1 antagonists has been discovered and developed (Hipskind, P. A. et. al. *Annu. Rep. Med. Chem.* **1996**, 31, 1-10; Rudolf, K. et. al. *Eur. J. Pharmacol.* **1994**, 271, R11; Serradeil-Le Gal, C. et. al. *FEBS Lett.* **1995**, 362, 192; Wright, J. et. al. *Bioorg. Med. Chem. Lett.* **1996**, 6, 1809; Poindexter, G. S. et. al. United States Patent 5,668,151; Peterson, J. M. et. al. WO9614307 (1996)). However, despite claims of activity in rodent models of feeding, it is unclear if inhibition of a feeding response can be attributed to antagonism of the Y1 receptor.


Several landmark studies strongly suggest that an "atypical Y1" receptor and / or the Y5 receptor, rather than the classic Y1 receptor, is responsible for invoking NPY-stimulated food consumption in animals. It has been shown that the NPY fragment NPY2-36 is a potent inducer of feeding despite poor binding at the classic Y1 receptor (Stanley, B. G. et. al. *Peptides* **1992**, 13, 581). Conversely, a potent and selective Y1 agonist has been reported to be inactive at stimulating feeding in animals (Kirby, D. A. et. al. *J. Med. Chem.* **1995**, 38, 4579). More pertinent to the invention described herein, [D-Trp³²]NPY, a selective Y5 receptor activator has been reported to stimulate food intake when injected into the hypothalamus of rats (Gerald, C. et. al. *Nature* **1996**, 382, 168). Since [D-Trp³²]NPY appears to be a full agonist of the Y5 receptor with no appreciable Y1 activity, the Y5 receptor is hypothesized to be

responsible for the feeding response. Accordingly compounds that antagonize the Y5 receptor should be effective in inhibiting food intake, particularly that stimulated by NPY.

5 A variety of structurally diverse compounds that antagonize the Y5 receptor have been described in various publications. In PCT WO 97/19682, aryl sulfonamides and sulfamides derived from arylalkylamines are described as Y5 antagonists and are reported to reduce food consumption in animals. In PCT WO 97/20820, PCT WO 97/20822 and PCT WO 97/20823, sulfonamides
10 containing heterocyclic systems such as quinazolin-2,4-diazirines, are likewise claimed as Y5 antagonists and reported to reduce feeding. In PCT WO 99/10330, a series of heterocyclic ketones is claimed to be NPY Y5 antagonists. In PCT WO 99/01128, certain diarylimidazole derivatives are claimed as a new class of NPY specific ligands. In PCT WO 98/35944, a
15 series of α -alkoxy and α -thioalkoxyamides are claimed to be NPY Y5 receptor antagonists. In PCT WO 98/35957, a series of amide derivatives are claimed as selective neuropeptide Y receptor antagonists; however, these compounds are structurally different from the compounds of this invention. The amides and amines of this invention that are described herein are novel molecular
20 entities that may have binding motifs that are different from these and other Y5 ligands that have been disclosed in patent applications or publications.

SUMMARY OF THE INVENTION

25 The present invention is related to compounds of formula A



5 R_1 is independently selected from the group consisting of hydrogen;
 hydroxy; halo; C_{1-8} alkyl; substituted C_{1-8} alkyl wherein the substituent is
 selected from halo, such as chloro, bromo, fluoro and iodo; C_{1-8} alkoxy;
 substituted C_{1-8} alkoxy wherein the substituent is selected from halo,
 such as chloro, bromo, fluoro and iodo; trifluoroalkyl; C_{1-8} alkylthio and
 substituted C_{1-8} alkylthio wherein the substituent is selected from halo,
 such as chloro, bromo, fluoro and iodo, trifluoro C_{1-8} alkyl and C_{1-8} alkoxy;
 C_{3-8} cycloalkyl; C_{3-8} cycloalkoxy; nitro; amino; C_{1-8} alkylamino; C_{1-}
 10 $_8$ dialkylamino; C_{4-8} cycloalkylamino; cyano; carboxy; C_{1-5} alkoxycarbonyl;
 C_{1-5} alkylcarbonyloxy; formyl; carbamoyl; phenyl and substituted phenyl
 wherein the substituent is selected from halo, hydroxyl, nitro, amino and
 cyano;

15 n is 1-2

B_1 is hydrogen;

B_2 is hydrogen;

20 or B_1 and B_2 may be methylene and joined together form a five or six-
 membered ring;

m 0-3

25 R_2 is independently selected from the group consisting of hydrogen;
 hydroxy; C_{1-6} alkyl; C_{2-6} alkenyl; halo, such as fluoro and chloro; C_{3-}
 $_7$ cycloalkyl; phenyl; substituted phenyl wherein the substituent is
 selected from halo, C_{1-6} alkyl, C_{1-6} alkoxy, trifluoro C_{1-6} alkyl, cyano, nitro,
 amino, C_{1-6} alkylamino, and C_{1-6} dialkylamino; naphthyl; substituted
 30 naphthyl wherein the substituent is selected from halo, C_{1-6} alkyl, C_{1-}
 $_6$ alkoxy, trifluoro C_{1-6} alkyl, cyano, nitro, amino, C_{1-6} alkylamino, and C_{1-}
 $_6$ dialkylamino; phenoxy; substituted phenoxy wherein the substituent is
 selected from halo, C_{1-6} alkyl, C_{1-6} alkoxy, trifluoro C_{1-6} alkyl, cyano and


nitro; a heteroaryl group such as pyridyl, pyrimidyl, furyl, thienyl, and imidazolyl; substituted heteroaryl wherein the substituent is selected from C₁₋₆alkyl and halo; and heterocycloalkyl such as pyrrolidino or piperidino;

5


Y is methylene (-CH₂-) or carbonyl (C=O)

L is selected from the group consisting of

C₁₋₈alkylene; C₂₋₁₀alkenylene; C₂₋₁₀alkynylene; C₃₋₇cycloalkylene;
 10 C₃₋₇cycloalkylC₁₋₄alkylene;
 arylC₁₋₄alkylene;
 α-aminoC₄₋₇alkylene;




15 (N-methylene)piperidin-4-yl;




(N-methylene)piperazin-4-yl;


20




(N-methylene)pyrrolidin-3-yl;



25 (N-methylene)-4-acetyl-piperidin-4-yl;




and (N-methylene)piperidin-4,4-diyl;




5 Z is selected from the group consisting of:

aryl;




10

N-sulfonamido;




N-(aryl)sulfonamido;




15

arylamido;




arylureido;




20


arylacetamido;



(aryloxy)carbonylamino;




2,3-dihydro-2-oxo-1H-benzimidazol-1-yl;



5

and 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl;



10

The aryl group in each case may be substituted as shown.

R_3 is independently selected from the group consisting of C_{1-8} alkyl; substituted C_{1-8} alkyl wherein the substituent is selected from C_{1-8} alkoxy and halo; cycloalkyl; substituted cycloalkyl wherein the substituent is selected from C_{1-8} alkoxy and halo; naphthyl; substituted naphthyl wherein the substituent is selected from halo, nitro, amino and cyano; heteroaryl wherein the heteroaryl group is selected from pyridyl, pyrimidyl, furyl, thienyl and imidazolyl; and substituted heteroaryl wherein the substituent is selected from halo, nitro, amino and cyano;

15

20

5 R_4 is independently selected from the group consisting of hydrogen; C_{1-8} alkyl; substituted C_{1-8} alkyl wherein the substituent is selected from alkoxy and halo; hydroxy; halogen; cyano; nitro; amino; C_{1-8} alkylamino and C_{1-8} dialkylamino; C_{1-8} alkoxy; substituted C_{1-8} alkoxy wherein the substituent is halo; hydroxy; halogen; cyano, nitro; amino and C_{1-8} alkylamino and C_{1-8} dialkylamino;

10 R_5 is independently selected from the group consisting of hydrogen; C_{1-8} alkyl; C_{1-8} alkylcarbonyl; aroyl; carbamoyl; amidino; (C_{1-8} alkylamino)carbonyl; (arylamino)carbonyl and aryl(C_{1-8} alkylcarbonyl);

R_6 is independently selected from the group consisting of hydrogen and C_{1-8} alkyl;

15 p is 1-3;

q is 1-3;

20 and enantiomers, diastereomers, and pharmaceutically acceptable salts thereof,

provided that:

25 when L is C_{1-8} alkylene, C_{2-10} alkenylene, C_{2-10} alkynylene, C_{3-7} cycloalkylene, C_{3-7} cycloalkyl C_{1-4} alkylene, aryl C_{1-4} alkylene or α -aminoalkylene;

then Z is phenyl, N-sulfonamido or N-(aryl)sulfonamido;

30 when L is (N-methylene)piperazin-4-yl;

then Z is phenyl or naphthyl;

when L is (N-methylene)pyrrolidin-3-yl or (N-methylene)piperidin-4-yl;

then Z is N-sulfonamido, N-(aryl)sulfonamido, 2,3-dihydro-2-oxo-
1*H*-benzimidazol-1-yl; benzamido, phenylureido,
phenylacetamido or (phenoxy)carbonylamino;

when L is (N-methylene)-4-acetyl-piperidin-4-yl;

then Z is phenyl or naphthyl and Y is carbonyl;

when L is (N-methylene)piperidin-4,4-diyl;

then Z is 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl and Y is
carbonyl;

and when B₁ and B₂ are both methylene thus forming a six-membered
ring (an aminotetralin) and when L is selected from the group consisting
of C₁₋₈alkylene; C₂₋₁₀alkenylene; C₂₋₁₀alkynylene or arylC₁₋₄alkylene;

then Z cannot be N-sulfonamido, N-(aryl)sulfonamido or phenyl;

all enantiomers and diastereomers of compounds of formula A are part
of the present invention, as are pharmaceutically acceptable salts
thereof.

Preferred compounds among the compounds of this invention are those
wherein B₁ and B₂ form a six-membered ring and m = 1-3.

As used herein unless otherwise noted the terms "alkyl" and "alkoxy"
whether used alone or as part of a substituent group, include straight and
branched chains having 1-8 carbon atoms. For example, alkyl radicals include
methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, *t*-butyl, pentyl, 2-
methyl-3-butyl, 1-methylbutyl, 2-methylbutyl, neopentyl, hexyl, 1-methylpentyl,

3-methylpentyl. Alkoxy radicals are oxygen ethers formed from the previously described straight or branched chain alkyl groups. The term "aryl" is intended to include phenyl and naphthyl and aroyl is intended to include arylacyl. The term "acyl" is intended to include C₁₋₈alkylcarbonyl. The term "halo", unless
5 otherwise indicated, includes bromo, chloro, fluoro and iodo. The term "cycloalkyl" is intended to include cycloalkyl groups having 3-7 carbon atoms. With reference to substituents, the term "independently" means that when more than one of such substituent is possible, such substituents may be the same or different from each other.

10

Those compounds of the present invention which contain a basic moiety can be converted to the corresponding acid addition salts by techniques known to those skilled in the art. Suitable acids which can be employed for this purpose include hydrochloric, hydrobromic, hydriodic, perchloric, sulfuric, nitric,
15 phosphoric, acetic, propionic, glycolic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, benzoic, cinnamic, mandelic, methanesulfonic, p-toluenesulfonic, cyclohexanesulfamic, salicylic, 2-phenoxybenzoic, 2-acetoxybenzoic, or saccharin, and the like. In general, the acid addition salts can be prepared by reacting the free base of compounds of
20 formula A with the acid and isolating the salt.

Pharmaceutical compositions containing one or more of the compounds of the invention described herein as the active ingredient can be prepared by intimately mixing the compound or compounds with a pharmaceutical carrier
25 according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral). Thus for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers,
30 coloring agents and the like; for solid oral preparations, such as powders, capsules and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Solid oral preparations may also be coated with substances such as

sugars or be enteric-coated so as to modulate the major site of absorption. For parenteral administration, the carrier will usually consist of sterile water and other ingredients may be added to increase solubility or preservation. Injectable suspensions or solutions may also be prepared utilizing aqueous carriers along with appropriate additives.

For the treatment of disorders of the central nervous system, the pharmaceutical compositions described herein will typically contain from 1 to about 1000 mg of the active ingredient per dosage; one or more doses per day may be administered. Determination of optimum doses and frequency of dosing for a particular disease state or disorder is within the experimental capabilities of those knowledgeable in the treatment of central nervous system disorders. The preferred dose range is 1-100 mg/kg.

As modulators of the NPY5 receptor, the compounds of Formula A are useful for treating feeding disorders such as obesity, anorexia nervosa and bulimia nervosa, and abnormal conditions such as epilepsy, depression, anxiety and sexual / reproductive disorders in which modulation of the NPY5 receptor may be useful. The compounds compete with the endogenous ligands NPY and PYY and possibly non-endogenous ligands, and bind to the NPY5 receptor. In addition, the compounds demonstrate antagonist activity by antagonizing the action of NPY upon binding to the Y5 receptor.


The compounds described herein are ligands of the NPY5 receptor, but are not necessarily limited solely in their pharmacological or biological action due to binding to this or any neuropeptide, neurotransmitter or G-protein coupled receptor. For example, the described compounds may also undergo binding to dopamine or serotonin receptors. The compounds described herein are potentially useful in the regulation of metabolic and endocrine functions, particularly those associated with feeding, and as such, may be useful for the treatment of obesity. In addition, the compounds described herein are potentially useful for modulating other endocrine functions, particularly those controlled by the pituitary and hypothalamic glands, and therefore may be


useful for the treatment of inovulation/infertility due to insufficient release of luteinizing hormone (LH) or luteal phase defect.


5 The present invention comprises pharmaceutical compositions containing one or more of the compounds of Formula A. In addition, the present invention comprises intermediates used in the manufacture of compounds of Formula A.


Examples of particularly preferred compounds of formula A include:


10




























DETAILED DESCRIPTION OF THE INVENTION

The amines and amides of formula **A** that comprise this invention are
5 synthesized via several distinct chemical syntheses as outlined in Schemes 1-
26; each synthetic route consists of several sequential chemical operations
that can be generalized as described below. In cases in which B₁ and B₂
together form a six-membered ring or a five-membered ring (an aminotetralin
or an aminoindane, respectively), the general synthesis entails the following
10 operations:

- Introduction of the α -substituent onto the tetralone (or indanone) nucleus
- Conversion to the corresponding α -substituted- β -aminotetralin (or α -
substituted-aminoindane)
- 15 • Acylation of the aminotetralin (or aminoindane) to afford amides of formula
A
- Reduction to produce amines of formula **A**

Protecting group manipulations may be needed at various stages of the
20 syntheses.

In cases where B₁ and B₂ are hydrogen, the general synthesis consists of the
following operations:

- 25 • Introduction of the α -substituent onto a phenylacetonitrile
- Reduction to the corresponding β -substituted phenethylamine
- Acylation of the phenethylamine to afford amides of formula **A**
- Reduction to produce amines of formula **A**


30 Protecting group manipulations may be needed at various stages of the
syntheses.

It is generally preferred that the respective product of each process step be separated from other components of the reaction mixture and subjected to purification before its use as a starting material in a subsequent step. Separation techniques typically include evaporation, extraction, precipitation and filtration. Purification techniques typically include column chromatography (Still, W. C. et. al., *J. Org. Chem.* **1978**, *43*, 2921), thin-layer chromatography, crystallization and distillation. The structures of the final products, intermediates and starting materials are confirmed by spectroscopic, spectrometric and analytical methods including nuclear magnetic resonance (NMR), mass spectrometry (MS) and liquid chromatography (HPLC). In the descriptions for the preparation of compounds of this invention, ethyl ether, tetrahydrofuran and dioxane are common examples of an ethereal solvent; benzene, toluene, hexanes and cyclohexane are typical hydrocarbon solvents and dichloromethane and dichloroethane are representative halohydrocarbon solvents. In those cases wherein the product is isolated as the acid addition salt the free base may be obtained by techniques known to those skilled in the art. In those cases in which the product is isolated as an acid addition salt, the salt may contain one or more equivalents of the acid.

Specifically, an appropriately substituted β -tetralone (II) is reacted with an aryl or heteroaryl aldehyde in the presence of a base such as piperidine, in an inert halohydrocarbon, ethereal or hydrocarbon solvent, such as benzene, from ambient temperature to reflux, to afford the corresponding α -benzylidenyl- β -tetralone or α -heteroarylmethylidenyl- β -tetralone (III). The β -tetralone (III) is dissolved in an inert hydrocarbon, ethereal, ester or alcohol solvent, such as methanol, and reacted with hydrogen gas at a pressure from ambient pressure to 100 p.s.i. in the presence of a suitable catalyst such as palladium on carbon. The reaction is performed at a temperature from ambient temperature to reflux, to yield the desired α -substituted- β -tetralone (IV) (Scheme 1).

An alternative method for the preparation of α -substituted- β -tetralones (IV) involves the reaction of an appropriately substituted β -tetralone (II) with a base such as pyrrolidine in an inert halohydrocarbon solvent such as

dichloromethane or hydrocarbon solvent such as benzene, under Dean-Stark conditions (removal of water) or in an alcohol solvent such as methanol, from ambient temperature to reflux, to afford enamine (V). Alkylation of enamine (V) is accomplished by reaction with a benzylic, heterocyclicalkyl or an allylic halide in an inert solvent such as acetonitrile, at a temperature from ambient temperature to reflux, to afford the α -substituted- β -iminium salt (VI). Hydrolysis of the salt (VI) to produce the desired α -substituted- β -tetralone product (IV) is accomplished by reaction of (VI) with water and an inorganic or organic acid such as hydrochloric or glacial acetic acid in an inert hydrocarbon, ethereal, alcohol or halohydrocarbon solvent, or a mixture thereof, such as methanol and dichloromethane (Scheme 1).




Scheme 1

15

The α -substituted- β -tetralones (IV) are converted to the corresponding aminotetralins via reaction with an ammonium salt such as ammonium acetate in the presence of a reducing agent such as sodium cyanoborohydride, for example, in an inert halohydrocarbon, hydrocarbon, ethereal or alcohol solvent such as methanol to produce the *cis*-aminotetralin (VII). In some cases, the *trans*-aminotetralin (VIII) is also formed as a minor product; both sets of diastereomers are part of this invention. The aminotetralins (VII) can also be

20

isolated as acid addition salts by treatment with an organic or an inorganic acid, such as trifluoroacetic acid or hydrochloric acid, for example (Scheme 2).




5

Scheme 2

Compounds in which $m = 0$ are prepared from an appropriately substituted aminotetralin (VII; $m = 0$) starting from 1-tetralones using the synthetic sequence shown in Scheme 2a.

10




Scheme 2a

Substituted phenethylamines (XI) are prepared by reacting an appropriately substituted phenylacetonitrile (IX) with an aryl or heteroaryl aldehyde in the presence of a base, such as sodium methoxide, in an inert alcohol solvent, such as methanol, at a temperature from ambient temperature to reflux, to afford α,β -unsaturated nitrile (X). Subsequent reduction of nitrile (X), for example, via reaction with hydrogen gas in the presence of a platinum oxide catalyst at a pressure from atmospheric pressure to approximately 100

20

psi, in an inert solvent such as aqueous alcohol, at a temperature from ambient temperature to reflux, affords β -substituted phenethylamine (XI). Alternatively, reaction of phenylacetonitrile (X) with an arylalkyl-, heteroarylalkyl- or alkyl halide, for example, such as allyl bromide in the presence of a base such as sodium methoxide or sodium hydride, in an inert solvent such as tetrahydrofuran or acetonitrile respectively, at a temperature from ambient to reflux, affords α -substituted phenylacetonitrile (XII). Subsequent reduction of nitrile (XII), for example, by hydrogenolysis, produces β -substituted phenethylamine (XI) (Scheme 3).

10



Scheme 3


The β -aminotetralins (VII) and the phenethylamines (XI) described above are acylated via suitable amidation methods (see Gross and Meienhofer, Eds., "The Peptides", Vols. 1-3, Academic Press, New York, NY, 1979-1981). A carboxylic acid is converted to an activated ester via peptide coupling methods known to those skilled in the art, and subsequently reacted with an aminotetralin (VII) or phenethylamine (XI), to afford the corresponding amides.

20

For example, a carboxylic acid such as *trans*-4-(2-fluorobenzenesulfonamido)methylcyclohexane carboxylic acid or 4-(*tert*-butoxycarbonyl)aminomethylcyclohexane carboxylic acid is reacted with HBTU


(2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate and an appropriate phenethylamine (XI), in the presence of a base such as diisopropylethylamine, in an inert solvent such as N,N-dimethylformamide, at a temperature from ambient temperature to reflux, to afford amide (XIII) or amide (XIV) respectively. Cleavage of the BOC (butoxycarbonyl) protecting group from carbamate (XIV) with trifluoroacetic acid produces the free amine, which is sulfonylated to yield amide (XIII).

The N-substituted phenethylamine compounds A of the invention are prepared via reduction of amide (XIII) by reaction with a suitable reducing agent such as borane-tetrahydrofuran complex or lithium aluminum hydride in an inert hydrocarbon solvent such as toluene or ethereal solvent such as tetrahydrofuran, at a temperature from ambient temperature to reflux. The final product can be isolated as an acid addition salt upon treatment with a suitable organic acid such as trifluoroacetic acid or an inorganic acid such as hydrochloric acid (Scheme 4).




Scheme 4

- 5 Aminotetralin analogs (B_1 and B_2 each are methylene) are prepared using the chemistry described above but replacing the phenethylamine (XI) starting material with an aminotetralin (VII) (Scheme 5).




Scheme 5

Compounds of formula A in which Z = 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl and L = (N-methylene)piperidin-4-yl are prepared from β -aminotetralins (VII) or phenethylamines (XI) and [4-(2-keto-1-benzimidazolyl)piperidin-1-yl]acetic acid (Schemes 6-7). For example, 4-(2-keto-1-benzimidazolyl)piperidine is reacted with a bromoacetic acid ester, such as ethyl bromoacetate, in the presence of an amine base, such as diisopropylethylamine, in an inert solvent such as acetonitrile, at a temperature ranging from ambient temperature to reflux, to afford ethyl [4-(2-keto-1-benzimidazolyl)piperidin-1-yl]acetate. This ester is subjected to hydrolysis under basic conditions, for example, by treatment with sodium hydroxide in an alcoholic solution such as aqueous methanol, to yield, upon acidification with an inorganic or organic acid such as hydrochloric or acetic acid for example, [4-(2-keto-1-benzimidazolyl)piperidin-1-yl]acetic acid. This carboxylic acid is reacted directly with β -aminotetralins (VII) or phenethylamines (XI), in the presence of an amine base, under peptide coupling conditions described above, to afford benzimidazolinones (XVII) and (XVIII) of formula A in which Y = carbonyl and L = (N-methylene)piperidin-4-yl (Schemes 6-7).




Scheme 6




Scheme 7

Compounds of formula A in which Y = methylene and L = (N-methylene)piperidin-4-yl and Z = 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl are prepared by reduction of amide (XVII) and amide (XVIII) with a reducing agent such as borane-tetrahydrofuran complex or lithium aluminum hydride as described above. The use of an aminotetralin (VII) starting material gives rise to products (XIX) (Scheme 8) whereas phenethylamines give the analogous amines (XX) (Scheme 9).




Scheme 8




Scheme 9

Compounds of formula A in which Y = carbonyl, L = (N-methylene)piperazin-4-yl and Z = phenyl are prepared by reacting a phenylpiperazine with a haloacetic acid ester, such as, for example, ethyl bromoacetate, in the presence of an amine base, such as diisopropylethylamine, in an inert solvent such as acetonitrile, at a temperature ranging from ambient temperature to reflux, to afford ethyl (4-aryl-piperazin-1-yl)acetate. This ester is subjected to hydrolysis under basic conditions, for example, by treatment with sodium hydroxide in an aqueous methanol, to yield, upon acidification with an inorganic or organic acid such as hydrochloric or acetic acid for example, (4-aryl-piperazin-1-yl)acetic acid. This carboxylic acid is reacted with β -aminotetralins (VII) or phenethylamines (XI), in the presence of a base, such as triethylamine for example, under peptide coupling conditions described above, to afford arylpiperidines (XXI) and (XXII) respectively, of formula A in which Y = carbonyl, L = (N-methylene)piperazin-4-yl and Z = aryl or substituted aryl (Schemes 10-11).




Scheme 10




Scheme 11

Compounds of formula **A** in which Y = methylene, L = (N-methylene)piperazin-4-yl and Z = aryl are prepared by reduction of amides (XXI) and (XXII) with a reducing agent such as borane-tetrahydrofuran complex or lithium aluminum hydride (see Scheme 9) to afford aminotetralins (XXIII) and phenethylamines (XIV) respectively (Schemes 12-13).



Scheme 12




Scheme 13


5

Replacement of 4-arylpiperazines with 4-arylpiperidines in Schemes 10 and 11 affords tetralinamides (XXV) and phenethylamides (XXVI) of formula A in which L = (N-methylene)piperidin-4-yl, Z = aryl and Y = carbonyl (Schemes 14-15).

10




Scheme 14



Scheme 15

Separately, reduction of amides (XXV) and (XXVI) with a reducing agent such a borane-tetrahydrofuran complex, affords amines (XXVII) and (XXVIII) of

formula A in which L = (N-methylene)piperidin-4-yl, Z = aryl and Y = methylene (Scheme 16).



Scheme 16


5

Compounds of formula A in which Y = carbonyl, L = (N-methylene)pyrrolidin-3-yl and Z = N-(aryl)sulfonamido are prepared by reacting a suitably protected aminopyrrolidine, such as (3-*t*-butoxycarbonylamino)pyrrolidine with a haloacetic acid ester, such as, for example, ethyl bromoacetate, in the presence of an amine base, such as diisopropylethylamine, in an inert solvent such as acetonitrile, at a temperature ranging from ambient temperature to reflux, to afford ethyl [(3-*t*-butoxycarbonylamino)pyrrolidin-1-yl]acetate. This ester is subjected to hydrolysis under basic conditions, for example, by treatment with sodium hydroxide in an aqueous methanol, to yield, upon acidification with an inorganic or organic acid such as hydrochloric or acetic acid for example, [(3-*t*-butoxycarbonylamino)pyrrolidin-1-yl]acetic acid. This carboxylic acid is reacted with β-aminotetralins (VII) or phenethylamines (XI), in the presence of a base, such as triethylamine for example, under peptide coupling conditions described above, to afford tetralinamides (XXIX) and phenethylamides (XXX) respectively. Subsequent treatment with an organic or inorganic acid, such as


20

trifluoroacetic acid and hydrochloric acid for example, produces the free terminal amines (XXXI) and (XXXII). These materials are sulfonylated by reaction with sulfonyl halides such as benzenesulfonyl chloride for example, in the presence of a base, to afford tetralinamides (XXXIII) and phenethylamides

5 (XXXIV) (Schemes 17-18).




Scheme 17




Scheme 18

Separately, reduction of amides (XXXIII) and (XXXIV) with a reducing
 5 agent such a borane-tetrahydrofuran complex, affords amines (XXXV) and
 (XXXVI) of formula A in which L = N-(methylene)pyrrolidin-3-yl and Z =
 sulfonamido or (aryl)sulfonamido, Y = methylene (Scheme 19).




Scheme 19

- 5 Tetralinamides and phenethylamides of formula A in which Y = carbonyl, L = (N-methylene)pyrrolidin-3-yl and Z = benzamido, phenylureido, phenylacetamido and phenoxycarbonylamino (or butoxycarbonylamino) are prepared by reacting amines (XXXI) and (XXXII) respectively, in an inert solvent at a temperature from ambient temperature to reflux, in the presence of
- 10 a base such as an amine or hydroxide, with an aroyl halide, an arylisocyanate, an arylacetyl halide or a chloroformate such as phenylchloroformate (or di-*tert*-butyl dicarbonate) to afford benzamides (XXXVII) and (XXXXI), phenylureas (XXXVIII) and (XXXXII), phenylacetamides (XXXIX) and (XXXXIII) and phenylcarbamate (XXXXX) and (XXXIV) respectively (Schemes 20-21).
- 15




Scheme 20




Scheme 21


Compounds of formula A in which Y = methylene, L = N-(methylene)pyrrolidin-3-yl and Z = benzamido, phenylureido, phenylacetamido and phenylcarbonylamino (or butoxycarbonylamino) are prepared by reducing amides (XXXI) and (XXXII) to their respective amines (XXXXV) and (XXXXVI) by treatment with a reducing agent such as borane-tetrahydrofuran complex or lithium aluminum hydride. Amines (XXXXV) and (XXXXVI) are subsequently separately reacted with an aroyl halide, an arylisocyanate, an arylacetyl halide or an arylchloroformate (or carbonate such as di-*tert*-butyl carbonate), in the presence of a base in an inert solvent as described in Scheme 20-21, to afford benzamides (XXXXVII) and (XXXXXI), phenylureas (XXXXVIII) and (XXXXXII), phenylacetamides (XXXXIX) and (XXXXXIII) and phenylcarbamates (XXXXX) and (XXXXXIV), respectively (Schemes 22-24).



Scheme 22



Scheme 23




Scheme 24

- 5 Substituting an appropriately protected aminopiperidine, such as (4-*t*-butoxycarbonylamino)piperidine for (3-*t*-butoxycarbonylamino)pyrrolidine in Schemes 17-24 affords compounds of formula A in which L = (N-methylene)piperidin-4-yl, Y = methylene or carbonyl and Z = N-(aryl)sulfonamido, sulfonamido, benzamido, phenylureido, phenylacetamido or
- 10 (phenoxy)carbonylamino.


Compounds of formula A in which Y = carbonyl, L = (N-methylene)piperidin-4,4-diyl and Z = 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl are prepared by reacting 1-aryl-1,3,8-triazaspiro-[4,5]decan-4-one with a

15 haloacetic acid ester, such as ethyl bromoacetate, in the presence of an amine

base, such as diisopropylethylamine, in an inert solvent such as acetonitrile, at a temperature from ambient temperature to reflux, to afford ethyl (1-aryl-1,3,8-triazaspiro-[4,5]decan-4-one-8-yl)acetate. This ester is subjected to hydrolysis under basic conditions, for example, by treatment with sodium hydroxide in an alcoholic solution such as aqueous methanol, to yield upon acidification with an inorganic or organic acid such as hydrochloric or acetic acid for example, (1-aryl-1,3,8-triazaspiro-[4,5]decan-4-one-8-yl)acetic acid. This carboxylic acid is reacted directly with β -tetralins (VII) or phenethylamines (XI), in the presence of a base such as triethylamine for example, under peptide coupling conditions described above, to afford aminotetralinamides (XXXXXV) and phenethylamides (XXXXXVI) respectively, of formula A in which Y = carbonyl, L = (N-methylene)piperidin-4,4-diyl and Z = 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl (Schemes 25-26).



Scheme 25




Scheme 26

5


Compounds of formula A in which L = (N-methylene)-4-acetyl-piperidin-4-yl and Z = phenyl are prepared by reacting 4-acetyl-4-phenylpiperidine with a haloacetic acid ester, such as, for example, ethyl bromoacetate, in the presence of an amine base, such as diisopropylethylamine, in an inert solvent such as acetonitrile, at a temperature ranging from ambient temperature to reflux, to afford ethyl [(4-acetyl-4-phenylpiperidin-1-yl)acetate]. This ester is subjected to hydrolysis under basic conditions, for example, by treatment with sodium hydroxide in an aqueous methanol, to yield, upon acidification with an inorganic or organic acid such as hydrochloric or acetic acid for example, [(4-acetyl-4-phenylpiperidin-1-yl)acetic acid]. This carboxylic acid is reacted with β -aminotetralins (VII) or phenethylamines (XI), in the presence of a base, such as triethylamine for example, under peptide coupling conditions described above, to afford (tetralinamido)arylpiperidines (XXXXXVII) and (phenethylamido)arylpiperidines (XXXXXVIII) respectively, of formula A in which Y = carbonyl, L = (N-methylene)-4-acetyl-piperidin-4-yl and Z = phenyl (Schemes 27-28).

20



Scheme 27

5



Scheme 28

Other compounds of this invention having the formula A can be prepared using the methods described herein; modifications of the experimental protocols described above are known or obvious or within the ability of those skilled in the art. For example, a variety of β -tetralones are known or readily prepared by reaction of phenylacetic acids with ethylene gas in the presence of a Lewis acid (for example, Stjernlof, P. et. al. *J. Med. Chem.* **1995**, *38*, 2202); these compounds can be directly converted to aminotetralins (VII) via reductive amination (Scheme 2). Phenethylamine intermediates (XI) are accessible from phenylacetone nitriles using literature methods (Jounral, Hawes and Wibberley, *J. Chem. Soc. C.* **1966**, 315 and 320; also see *J. Am. Chem. Soc.* **1989**, *111*, 5954 and *Synthesis* **1997**, *11*, 1268) and can be used to prepare compounds of formula A in which B₁ and B₂ are both hydrogen (Scheme 3). Compounds in which the R₁ group(s) is varied can be obtained using the chemistry described above; in some cases, protecting group manipulations are used and these are obvious or known to those skilled in the art. Examples include masking an amine group as a carbamate, amide or phthalamide, and masking an hydroxyl group as an ether or ester. Other R₁ substituents are available through functional group manipulations such as, for example, reduction of a nitro group to an amine or dehydration of an amide to a nitrile.

Variation of the R₂ group is readily accomplished by using substituted benzaldehydes, naphthylaldehydes and heteroaryl carboxaldehydes, or by using alkyl, alkylenic, alkynyl and benzylic halides, or by using phenoxyalkyl and haloalkyl halides in Schemes 1 and 3. Compounds in which the L group is varied, are derived from piperazines, piperidines or pyrrolidines as described in Schemes 6, 10, 14, 17 and 25. Compounds in which L is alkylene, alkenylene, alkynylene, cycloalkylene or cycloalkylalkylene are derived from amino-carboxylic acids such as aminohexanoic acid, aminohexenoic acid, aminohexynoic acid. Compounds in which L is α -aminoalkylene are derived

from amino acids such as lysine which can be used in the racemic or enantiomeric form.

Compounds of formula **A** where Z is sulfonamido or (aryl)sulfonamido, in which either the R₃ or the R₄ group is varied, are accessible by sulfonylation; there are hundreds of sulfonyl halides or sulfonic acids that are commercially available and more that are known. Compounds of formula **A** where Z is sulfonamido or (aryl)sulfonamido, in which the R₃ substituent is heteroaryl can be prepared by substituting a pyridinyl, thienyl or furyl sulfonylchloride for a benzenesulfonamide as described in Schemes 4-5. Similarly, alkylsulfonyl and cycloalkylsulfonyl halides, alone or in the presence of an activating agent such as a Lewis acid, can be used to prepare sulfonamides of formula **A** in which the R₃ substituent is alkyl or cycloalkyl respectively. Compounds in which Z is phenyl or aryl are obtained directly from arylpiperazines and arylpiperidines as described in Schemes 10 and 14 respectively; hundreds of arylpiperazines and arylpiperidines are known or commercially available and can be used to make compounds of this invention. Compounds of formula **A** where Z is benzamido, phenylureido, phenylacetamido, (phenoxy)carbonylamino are prepared from aroyl halides, isocyanates, phenylacetyl halides and chloroformates as described in Schemes 20-21 and 23-24 and hundreds of reagents of these kinds are commercially available or known.

Compounds of formula **A** in which B₁ and B₂ are joined together to form a five-membered ring (an aminoindane) are prepared starting from an indanone and using the chemistry described herein. It is preferable to use a symmetrical indan-2-one to avoid the formation of regiochemical isomers which are difficult to separate.

EXAMPLES

The following examples describe the invention in greater detail and are intended to illustrate the invention, but not to limit it. All compounds were identified by a variety of methods including nuclear magnetic resonance spectroscopy, mass spectrometry and, in some cases, infrared spectroscopy and elemental analysis. Nuclear magnetic resonance (300 MHz NMR) data are reported in parts per million downfield from tetramethylsilane. Mass spectra data are reported in mass/charge (m/z) units. Unless otherwise noted, the materials used in the examples were obtained from readily available commercial sources or synthesized by standard methods known to those skilled in the art.

EXAMPLES 1-2

2-Amino-6-[(2-fluorophenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthenyl-(2*S*)-hexanamide bis-hydrochloride 7

N-[5-amino-6-[[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenyl]amino]hexyl-2-fluorobenzenesulfonamide tris-hydrochloride 8

A. 6-Methoxy- β -tetralone 1 (2.0 g, 11.3 mmol) and diisopropylethylamine (0.20 mL, 1.1 mmol) were dissolved in benzene (60 mL) with stirring in a 100 mL round-bottom flask. 3-Pyridylcarboxaldehyde (1.1 mL, 11.7 mmol) was added and the reaction vessel was flushed with argon and a Dean-Stark trap with reflux condenser was attached. The mixture was heated at reflux for 19 hours. After cooling, HPLC analysis indicated that no products had formed. Piperidine (0.094 mL, 1.1 mmol) was added at this time and heating at reflux was continued for 23 hours. The solvents were removed *in vacuo* to yield a glassy orange solid. Chromatographic purification [silica gel column (dimensions 5 x 29 cm) eluting with a gradient of: 100% hexane (400 mL),

75%/25% hexane/ethyl acetate (v/v) (400 mL), 50%/50% hexane/ethyl acetate (v/v) (400 mL), 25%/75% hexane/ethyl acetate (v/v) (400 mL), and finally with 100% ethyl acetate] was performed. After evaporation of the appropriate fractions, 3,4-dihydro-6-methoxy-1-((3-pyridinyl)methylidenyl)-2-naphthalenone 2 (1.484 g, 5.59 mmol) was obtained as an orange oil which solidified upon standing in the refrigerator. MS (MH⁺) 266; ¹H NMR (CDCl₃) δ 2.67 (t, 2H), 3.02 (t, 2H), 3.83 (s, 3H), 6.60 (dd, 1H), 6.82 (d, 1H), 7.19 (m, 2H), 7.51 (s, 1H), 7.71 (d, 1H), 8.49 (dd, 1H), 8.65 (d, 1H).

10 B. The naphthalen-2-one 2 (1.442 g, 5.44 mmol) obtained above was dissolved in absolute ethanol (50 mL) and transferred to a 250 mL Parr hydrogenation bottle. Separately, ethanol was carefully added to 10% palladium on carbon (0.020g) and this slurry was added to the Parr bottle. The mixture was hydrogenated under a pressure of 50 psi for 16 hours. The catalyst was removed by filtration over Celite. Spectroscopic evidence indicated the presence of some starting material and so more palladium catalyst (0.081 g) was added to the ethanol solution and the hydrogenation was repeated for 20 hours. The catalyst was then removed by filtration over Celite. Removal of the solvents *in vacuo* yielded 3,4-dihydro-6-methoxy-1-(3-pyridinylmethyl)-2(1H)-naphthalenone 3 as an orange oil which was used in the next step without further purification. MS (MH⁺) 268.

C. Naphthalen-2-one 3 obtained above was dissolved in methanol (275 mL) in a 1 L round-bottom flask. Ammonium acetate (4.27 g, 55.4 mmol) was added to the stirred methanol solution and was allowed to completely dissolve before proceeding. Sodium cyanoborohydride (1.703 g, 27.5 mmol) was then added to the methanol solution. The reaction vessel was flushed with nitrogen and the solution refluxed for 18 hours. The solvents were then removed *in vacuo* to yield a yellow solid which was dissolved in ethyl ether (500 mL) and 0.1 M sodium hydroxide solution (275 mL). The organic layer was removed and washed with an additional 0.1 M sodium hydroxide solution (275 mL) and with water (250 mL). The combined aqueous washes were back extracted with ethyl ether (3 x 100 mL). The organic extracts were combined and dried

over sodium sulfate. The solvents were removed *in vacuo* and the residue was taken up in ethyl ether and a minimum amount of dichloromethane. An excess of 1 M hydrogen chloride in ethyl ether was added and a dark tan precipitate formed. The solvents were removed *in vacuo* and the resulting solid was triturated with ether and dried in a vacuum oven to yield 1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenamine bis-hydrochloride **4** as a tan-orange solid (1.208 g, 3.54 mmol) MS (MH^+) 269; 1H NMR ($DMSO-d_6$) δ 1.95-2.20 (m, 2H), 2.68-3.29 (m, 4H), 3.30-3.48 (m, 2H), 3.69 (s, 3H), 5.98 (d, 1H), 6.41 (dd, 1H), 6.75 (d, 1H), 7.98 (dd, 1H), 8.36 (d, 1H), 8.68-8.89 (m, 5H) (Figure 1).




Figure 1

D. *N*-*tert*-Butoxycarbonyl-L-Lysine (2.49 g, 10.1 mmol) was placed in a 200 mL round-bottom flask. A magnetic stir bar was added followed by 10 mL dioxane and 21 mL 1N sodium hydroxide solution. The solution was stirred for several minutes until complete dissolution had occurred. A solution of 2-fluorobenzenesulfonyl chloride (2.00 g, 10.3 mmol) in dioxane (11 mL) was added via pipette. The reaction vessel was flushed with argon, capped and allowed to stir at ambient temperature for approximately 1.5 hours. The stir bar was then removed and the solvent evaporated under reduced pressure until

WO 01/09120

only water remained. To this mixture water was added to bring the volume to about 50 mL and 1N hydrochloric acid (22 mL) was added which resulted in the formation of a gooey precipitate. This mixture was extracted with methylene chloride (3 x 50 mL) and the combined organics were washed with 1N hydrochloric acid (1 x 50 mL) and then brine (1 x 50 mL). The organics were dried over magnesium sulfate, filtered and concentrated *in vacuo* to yield the sulfonated N-t-butoxycarbonyl-lysine 5 (3.93 g, 9.7 mmol) as an off-white glassy semi-solid. NMR(d_6 -DMSO): δ 12.42 (s, 1H), 7.90 (t, 1H), 7.79 (t, 1H), 7.71 (m, 1H), 7.49-7.34 (m, 2H), 7.02 (d, 1H), 3.78 (m, 1H), 2.83 (m, 2H), 1.63-1.16 (m, 15H); MS: M-H = 403.

E. The sulfonated L-lysine 5 from the previous reaction (3.92 g, 9.69 mmol) was placed in a 300 mL round-bottom flask along with 1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenamine bis-hydrochloride 4 (3.53 g, 10.34 mmol) and a stir bar. N,N-Dimethylformamide (DMF) (50 mL) was added followed by diisopropylethylamine (5.6 mL, 32.1 mmol) and the mixture was stirred. After dissolution, 2-(1H-benzotriazole-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate (HBTU) (3.72 g, 9.81 mmol) was added and the flask was flushed with argon, capped and allowed to stir at ambient temperature for 30 minutes. Water (~5 mL) was then added to quench the reaction and the solvents were removed *in vacuo* to give a brown oil. This material was purified by column chromatography on a silica gel column (dimensions 6 x 12 cm) eluting with a gradient of methylene chloride-acetone-methanol. After evaporation of the appropriate fractions, adduct 6 (as a tan-green foam, 4.63 g, 7.07 mmol) was obtained as a mixture of diastereomers. MS: MH^+ = 655.

F. The sulfonated lysino-tetralinamide 6 from the previous reaction (4.59 g, 7.01 mmol) was placed in a 200 mL round-bottom flask with a stir bar and methylene chloride (100 mL) was added. With stirring, a solution of 95%TFA / 5% H₂O (v/v) (10 mL) was added and the reaction mixture was allowed to stir

under nitrogen at ambient temperature for 3.5 hours. The reaction mixture was then concentrated *in vacuo* and the residue was triturated with diethyl ether. The liquid was decanted and more ether was added. The resultant solid was filtered and dried under vacuum to give the desired tetralinamide lysino-

5 sulfonamide bis-hydrochloride **7** (4.28 g, 5.47 mmol) as a mixture of diastereomers. A portion of this material (4.01 g) was separated into racemic sets of diastereomers via reverse-phase chromatography (Bondapak C18, 6x(40x100mm) column using a gradient of H₂O/CH₃CN (+0.1%TFA)). The appropriate fractions were isolated and lyophilized to yield diastereomer **a**

10 (2.17 g, 2.77 mmol) and diastereomers **b** (1.78 g, 2.27 mmol) as bis-TFA salts (absolute configurations of the diastereomers were not determined). Diastereomer **a**: de = 96%; NMR(d₆-DMSO): δ 8.57 (m, 2H), 8.30 (s, 1H), 8.11 (br, 3H), 7.96 (t, 1H), 7.80-7.64 (m, 3H), 7.55 (dd, 1H), 7.48-7.32 (m, 2H), 6.71 (s, 1H), 6.58-6.46 (m, 2H), 4.03 (m, 1H), 3.79 (m, 1H), 3.69 (s, 3H), 3.24 (m,

15 1H), 3.03-2.73 (m, 6H), 2.08-1.91 (m, 1H), 1.85-1.58 (m, 3H), 1.53-1.31 (m, 4H); MS: MH⁺ = 555. Diastereomer **b**: de = 100%; NMR(d₆-DMSO): δ 8.68 (d, 1H), 8.57 (d, 1H), 8.49 (s, 1H), 8.21 (br, 3H), 8.01 (d, 1H), 7.93 (t, 1H), 7.78 (dt, 1H), 7.73 (m, 2H), 7.52-7.37 (m, 2H), 6.75 (s, 1H), 6.56 (m, 2H), 3.99 (m, 1H), 3.85 (m, 1H), 3.71 (s, 3H), 3.23 (m, 1H), 3.08-2.76 (m, 6H), 2.00-1.59 (m, 4H),

20 1.53-1.22 (m, 4H); MS: MH⁺ = 555 (Figure 2).

G. Diastereomer **a** **7** from the previous reaction (2.02 g, 2.58 mmol) was placed in a 200 mL round-bottom flask along with a stir bar and THF (60 mL) was added. After stirring, a solution of borane in THF (40 mL of a 1M solution,

25 40 mmol) was added and the flask was flushed with nitrogen and a reflux condenser was attached. The mixture was heated at reflux for 24 hours at which time an additional portion of the borane solution (10 mL) was added. The reaction mixture was heated at reflux for an additional 14 hours. The reaction mixture was allowed to cool and water (10 mL) was carefully added to

30 quench the reaction. Hydrochloric acid (20 mL of a 1N solution) was added and the reaction mixture was heated at reflux for 2 hours. The solvents were removed *in vacuo* and the residue was suspended in water (250 mL). This

- mixture was made slightly acidic via the addition of 1N hydrochloric acid. This aqueous solution was washed with methylene chloride (3 x 250 mL) and the aqueous layer was separated. Ammonium hydroxide solution was added until the pH was basic. The water was then removed *in vacuo* giving a white solid.
- 5 The resultant material was triturated with methylene chloride and the borane salts that precipitated were removed by filtration. The remaining organics were concentrated *in vacuo* to give the crude product as a foam. This material was purified by flash chromatography on a silica gel column (dimensions 6 x 11 cm) eluting with a gradient of methylene chloride-methanol-ammonium hydroxide.
- 10 After evaporation of the appropriate fractions, the residue was treated with an excess of ethanolic-hydrogen chloride, followed by evaporation and drying under vacuum, to obtain aminotetralin sulfonamide 8 as a yellow tris-hydrochloride salt (0.898 g, 1.38 mmol). NMR(d_6 -DMSO): δ 10.83 (br, 1H), 10.08 (br, 1H), 8.80 (d, 1H), 8.73 (m, 4H), 8.43 (d, 1H), 7.97 (m, 2H), 7.81 (t, 1H), 7.71 (m, 1H), 7.51-7.33 (m, 2H), 6.75 (s, 1H), 6.37 (d, 1H), 5.83 (d, 1H), 3.80 (m, 1H), 3.71-3.30 (m, 8H), 3.11 (m, 1H), 2.98-2.69 (m, 4H), 2.34-2.13 (m, 2H), 1.73-1.55 (m, 2H), 1.54-1.29 (m, 4H); MS: MH⁺ = 541 (Figure 2).
- 15




Figure 2

EXAMPLE 3

N-[5-amino-6-[[cis-1,2,3,4-tetrahydro-6-hydroxy-1-(3-pyridinylmethyl)-2-naphthalenyl]amino]hexyl-2-fluorobenzenesulfonamide tris-hydrochloride **9**

5

Aminotetralin sulfonamide **8** from the previous reaction (0.160 g, 0.246 mmol) was placed in a 50 mL round-bottom flask along with a stir bar. Methylene chloride (25 mL) was added and the slurry was cooled on an ice bath for several minutes. Boron tribromide in methylene chloride (1M, 1.25 mL, 1.25 mmol) was added to the reaction. The flask was flushed with argon, capped and allowed to warm up to ambient temperature and the mixture was stirred over 16 hours at which time the reaction was quenched by the addition of methanol (1 mL). The solvents were removed in vacuo and an additional aliquot of methanol was added to the resultant residue. Evaporation of the solvent from this mixture afforded crude product which was purified via reverse-phase chromatography (Bondapak C18, 3x(40x100mm), gradient of H₂O / CH₃CN (+0.1%TFA)). The appropriate fractions were collected and lyophilized. The resultant material was subsequently treated with ethanolic-hydrogen chloride, followed by evaporation and drying under vacuum to give the phenolic product **9** as a white tris-hydrochloride salt (0.145 g, 0.228 mmol). NMR(d₆-DMSO): δ 10.77 (br, 1H), 10.01 (br, 1H), 9.31 (br, 1H), 8.79 (d, 1H), 8.67 (m, 4H), 8.37 (d, 1H), 7.97 (m, 2H), 7.81 (dt, 1H), 7.72 (m, 1H), 7.52-7.36 (m, 2H), 6.57 (s, 1H), 6.22 (dd, 1H), 5.69 (d, 1H), 3.79 (m, 1H), 3.68-3.30 (m, 5H), 3.04 (m, 1H), 2.92-2.68 (m, 4H), 2.33-2.10 (m, 2H), 1.73-1.56 (m, 2H), 1.55-1.32 (m, 4H); MS: MH⁺ = 527 (Figure 3).

10

15

20

25




Figure 3

5 EXAMPLE 4

(2S)-2-(Acetylamino)-6-[(2-fluorophenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthenyl]hexanamide bis-
 10 hydrochloride 10

Diasteromerically mixed tetralinamide lysino-sulfonamide 7 (0.195 g, 0.249 mmol) was placed into a 50 mL round-bottom flask along with a stir bar. Acetonitrile (25 mL) was added followed by triethylamine (0.122 mL, 0.875 mmol). With stirring, acetyl chloride (0.021 mL, 0.295 mmol) was added and the flask was flushed with argon, capped and stirred overnight at ambient temperature. The solvents were removed *in vacuo* and the residue was taken up in methylene chloride (75 mL). This mixture was washed with 1N sodium hydroxide (2 x 25 mL) and then with brine (1 x 25 mL). The organics were dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the acetate product 10 as a tan solid (0.139 g, 0.233 mmol) as a 1:1
 20

diastereomeric mixture. NMR(CDCl₃): δ 8.52 (d, 0.5H), 8.43 (d, 0.5H), 8.28 (d, 1H), 7.89 (m, 1H), 7.57 (m, 1H), 7.44 (d, 0.5H), 7.39-7.13 (m, 3.5H), 6.92 (t, 0.5H), 6.77 (d, 0.5H), 6.70-6.54 (m, 3H), 6.48 (dd, 1H), 6.34 (d, 0.5H), 5.59 (t, 0.5H), 4.40-4.06 (m, 2H), 3.78 (d, 3H), 3.29 (m, 1H), 3.19-2.82 (m, 6H), 2.01 (d, 3H), 1.92-1.71 (m, 2H), 1.72-1.32 (m, 6H); MS: MH⁺ = 597 (Figure 4).

EXAMPLE 5

10 (2S)-2-(Acetylamino)-6-[(2-fluorophenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-hydroxy-1-(3-pyridinylmethyl)-2-naphthenyl]hexanamide bis-hydrochloride **11**

15 The bis-amide **10** from the previous reaction (0.114 g, 0.191 mmol) was placed in a 50 mL round-bottom flask along with a stir bar. Methylene chloride (20 mL) was added and the solution was cooled on an ice bath for several minutes. Boron tribromide in methylene chloride (1M, 1.0 mL, 1.0 mmol) was added to the reaction mixture. The flask was flushed with argon, capped and allowed to
20 warm up to ambient temperature and the mixture was stirred over 16 hours at which time the reaction was quenched by the addition of methanol (1 mL). The solvents were removed *in vacuo* and the resultant material treated with an additional aliquot of methanol. This mixture was evaporated *in vacuo* to yield crude phenolic tetralinamide **11** which was purified via reverse-phase column
25 chromatography which allowed for separation and purification of the racemic pairs of diastereomers (Bondapak C18, 3x(40x100mm), gradient of H₂O/CH₃CN (+0.1%TFA)). After lyophilization of the appropriate fractions, each diastereomer was treated with ethanolic-hydrogen chloride, subjected to evaporation and lastly dried under vacuum to give the individual racemic
30 diastereomers as tan hydrochloride salts; diastereomer a (0.036 g, 0.058 mmol) and diastereomer b (0.057 g, 0.092 mmol) (absolute configurations of the diastereomers were not determined). Diastereomer a: de = 100%; NMR(d₆-DMSO): δ 9.22 (v. br, 1H), 8.79 (d, 1H), 8.48 (s, 1H), 8.20 (d, 1H),

8.08-7.87 (m, 4H), 7.83-7.63 (m, 2H), 7.50-7.33 (m, 2H), 6.54 (s, 1H), 6.43-6.28 (m, 2H), 4.19 (q, 1H), 3.93 (m, 1H), 3.18 (m, 1H), 3.08-2.67 (m, 6H), 1.92 (m, 1H), 1.84 (s, 3H), 1.73 (m, 1H), 1.58-1.16 (m, 6H); MS: MH⁺ = 583. Diastereomer **b**: de = 66%; NMR(d₆-DMSO): δ 9.20 (v. br, 1H), 8.77 (d, 1H), 8.57 (s, 1H), 8.28-8.14 (m, 2H), 8.08-7.84 (m, 3H), 7.83-7.62 (m, 2H), 7.50-7.32 (m, 2H), 6.54 (s, 1H), 6.47-6.29 (m, 2H), 4.10 (q, 1H), 3.85 (m, 1H), 3.27-3.08 (m, 2H), 3.03-2.66 (m, 5H), 1.90 (s, 3H), 1.87-1.63 (m, 2H), 1.57-1.13 (m, 6H); MS: MH⁺ = 583 (Figure 4).

10




Figure 4

EXAMPLE 6

5 3-[(Phenylsulfonyl)amino]-N-[cis-1,2,3,4-tetrahydro-6-fluoro-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-pyrrolidineacetamide bis-trifluoroacetate 17

A. Racemic 3-(N-butoxycarbonyl)aminopyrrolidine (5.13 g, 27.5 mmol) was placed into a 300 mL round-bottom flask along with a stir bar. Acetonitrile (100 mL) was added which gave a slurry to which was added diisopropylethylamine (7.2 mL, 41.3 mmol) followed by ethyl bromoacetate (3.1 mL, 28.0 mmol). The flask was flushed with nitrogen and a reflux condenser was attached. The reaction mixture was heated at reflux for 1.5 hours then allowed to cool and stir at ambient temperature overnight. The solvents were removed *in vacuo* to give an oily solid. This material was taken up in methylene chloride (200 mL) and washed successively with sodium bicarbonate solution (1 x 200 mL), water (1 x 200 mL) and brine (200 mL). The organics were dried over magnesium sulfate, filtered and the solvents removed *in vacuo* to give a thick oil which slowly crystallized upon standing to give the pyrrolidinylacetate ester 12 (6.96 g, 25.6 mmol). NMR(CDCI₃): δ 4.98 (br d, 1H), 4.27-4.13 (m, 3H), 3.33 (s, 2H), 2.98 (m, 1H), 2.83-2.66 (m, 2H), 2.48 (m, 1H), 2.27 (m, 1H), 1.67 (m, 1H), 1.44 (s, 9H), 1.28 (t, 3H).

B. Pyrrolidinylacetate ester 12 from the previous reaction (6.95 g, 25.5 mmol) was put into a 300 mL round-bottom flask. A stir bar and methanol (100 mL) was added. The mixture was stirred until all of the starting material had dissolved. Sodium hydroxide solution (1N, 75.0 mL, 75.0 mmol) was added to the resulting solution. The reaction vessel was capped and the mixture was allowed to stir for 20 hours at which time hydrochloric acid was added (1N, 75.0 mL, 75.0 mmol). The resultant mixture was allowed to stir for several minutes. The solvents were removed *in vacuo* and the resulting solid was treated with methylene chloride. The organic extract was dried over magnesium sulfate, filtered and concentrated *in vacuo* to give pyrrolidinylacetic

acid **13** as a white powder (6.30 g, 25.8 mmol). NMR(d_6 -DMSO): δ 7.21 (br d, 1H), 4.05 (m, 1H), 3.38 (s, 2H), 3.23 (m, 1H), 3.02 (m, 2H), 2.78 (m, 1H), 2.12 (m, 1H), 1.73 (m, 1H), 1.39 (s, 9H); MS: $MH^+ = 245$.

- 5 C. 1,2,3,4-Tetrahydro-6-fluoro-1-(3-pyridinylmethyl)-2-naphthalenamine bis-hydrochloride **14** (0.331 g, 1.01 mmol), prepared from 6-fluoro- β -tetralone using the chemistry described in EXAMPLE 1 (Figure 1), was placed in a 25 mL round-bottom flask along with a stir bar and DMF (5 mL) was added. The pyrrolidinylacetic acid **13** (0.250 g, 1.02 mmol) from the previous reaction was
10 added followed by diisopropylethylamine (0.580 mL, 3.33 mmol) and then HBTU (0.387 g, 1.02 mmol). The flask was flushed with argon, capped and allowed to stir at ambient temperature for 2 hours. The reaction was diluted with brine (50 mL) and methylene chloride (150 mL) and the layers separated. The organics were washed with more brine (2 x 50 mL). The combined
15 aqueous brine washes were extracted with methylene chloride (2 x 25 mL) and the combined organics were dried over magnesium sulfate, filtered and concentrated *in vacuo* to give the crude product. This material was purified via reverse-phase column chromatography (Bondapak C18, 3x(40x100mm), gradient of H_2O/CH_3CN (+0.1%TFA)). Lyophilization of the appropriate
20 fractions gave the pyrrolidineacetamide bis-TFA salt **15** as a white powder (0.251 g, 0.35 mmol); MS: $MH^+ = 483$.

D. Pyrrolidineacetamide **15** from the previous reaction (0.205 g, 0.288 mmol) was placed in a 50 mL round-bottom flask along with a stir bar. Methylene chloride (25 mL) was added followed by a small amount of water (~0.5 mL) and TFA (2 mL). The reaction was capped and allowed to stir at ambient temperature for 19 hours at which time the solvents were removed *in vacuo* to yield 3-aminopyrrolidineacetamide tris-TFA salt **16** (0.204 g, 0.282 mmol). NMR(d_6 -DMSO): δ 8.69 (d, 1H), 8.64 (d, 1H), 8.49 (s, 1H), 8.36 (br, 3H), 7.93 (d, 1H), 7.67 (t, 1H), 7.02 (d, 1H), 6.83 (m, 2H), 4.13 (s, 2H), 4.07-3.88 (m, 3H), 3.87-3.22 (m, 4H), 3.15-2.69 (m, 4H), 2.41 (m, 1H), 2.14-1.69 (m, 3H); MS: MH⁺ = 383.

E. Aminopyrrolidine acetamide **16** from the previous reaction (0.074 g, 0.102 mmol) was placed into a 50 mL round-bottom flask along with a stir bar and acetonitrile (20 mL) was added. Diisopropylethylamine (0.078 mL, 0.448 mmol) was added followed by benzenesulfonyl chloride (0.013 mL, 0.102 mmol). The flask was flushed with argon, capped and allowed to stir at ambient temperature for 3 hours at which time the solvents were removed *in vacuo*. The residue was purified by reverse-phase column chromatography (H₂O/CH₃CN (+0.1%TFA)). After isolation and lyophilization of the appropriate fractions, 3-[(phenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-fluoro-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-pyrrolidineacetamide bis-TFA salt **17** was obtained as a white solid (0.067 g, 0.089 mmol). NMR(d_6 -DMSO): δ 8.62 (d, 2H), 8.47 (s, 1H), 8.25 (m, 1H), 7.92 (d, 1H), 7.83 (m, 2H), 7.66 (m, 4H), 7.02 (d, 1H), 6.84 (m, 2H), 4.18-3.73 (m, 4H), 3.72-2.72 (m, 9H), 2.07 (m, 1H), 1.98-1.67 (m, 3H); MS: MH⁺ = 523 (Figure 5).




Figure 5

EXAMPLES 7-8

4-(2,3-Dihydro-2-oxo-1*H*-benzimidazol-1-yl)-*N*-[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-
5 hydrochloride 19

4-(2,3-Dihydro-2-oxo-1*H*-benzimidazol-1-yl)-*N*-[*trans*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-
10 hydrochloride 20

A solution of 2-(1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) (0.974 g, 2.57 mmol), 4-(2,3-dihydro-2-oxo-1*H*-benzimidazol-1-yl)-1-piperidineacetic acid (1.20 g, 2.57 mmol), and *N,N*-diisopropylethylamine (1.8 mL, 10.3 mmol) in *N,N*-dimethylformamide (15 mL)
15 was stirred at room temperature for 5 min. To this mixture, 1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenamine bis-hydrochloride 4 (0.80 g, 2.34 mmol) was added and stirring was continued for 18 h. The solution was heated to 100°C for 1 h. The solution was cooled and poured into a saturated solution of aqueous sodium bicarbonate. A fine green precipitate
20 was collected by filtration, and the solid was purified by reverse phase C₁₈ HPLC eluted with a gradient of water/acetonitrile/trifluoroacetic acid 10/90/0.1 to 90/10/0.1. The *cis* product 19 was isolated as a colorless solid (0.386 g, 22%): ¹H NMR (DMSO-*d*₆) δ 1.76 (m, 4 H), 2.72-3.02 (m, 4 H), 3.16 (d, 2 H), 3.29-3.46 (m, 3 H), 3.54-3.75 (m, 2 H) superimposed on 3.72 (s, 3 H), 3.92-
25 4.07 (m, 3 H), 4.53-4.65 (m, 1 H), 6.63 (d, 1 H), 6.70-6.77 (m, 2 H), 7.04 (br s, 3 H), 7.59 (br s, 1 H), 7.99 (t, 1 H), 8.37 (d, 1 H), 8.74 (m, 2 H), 8.96 (d, 1 H), 10.5-10.71 (br s, 1 H), and 11.03 (s, 1 H); MS *m/e* 512 (MH⁺). A mixture of *cis/trans* isomers ~8/2 0.490 g (28%) was also obtained as well as the purified *trans* isomer 20 as a colorless solid (0.136 g, 8%): ¹H NMR (DMSO-*d*₆) δ 1.70
30 (m, 6 H), 2.63-3.81 (m, 9 H) superimposed on 3.72 (s, 3 H), 3.83-4.00 (m, 3 H), 4.47-4.60 (m, 1 H), 6.67-6.82 (m, 3 H), 7.02 (br s, 3 H), 7.21 (d, 1 H), 7.70 (t, 1 H), 8.14 (d, 1 H), 8.50-8.73 (m, 3 H), 9.70-10.10 (br s, 1 H), and 11.0 (s, 1 H); MS *m/e* 512 (MH⁺) (Figure 6).




Figure 6

EXAMPLE 9

4-Acetyl-4-phenyl-N-[cis-1,2,3,4-tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-hydrochloride **21**

- 5 1,2,3,4-Tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenamine bis-hydrochloride
4 (0.75 g, 2.41 mmol) was reacted with 2-(4-acetyl-4-phenyl-piperidin-1-yl)acetic acid (0.86 g, 2.65 mmol), *N,N*-diisopropylethylamine (2.0 mL, 11.3 mmol) and HBTU (1.01 g, 2.65 mmol) in *N,N*-dimethylformamide (15 mL) at
10 room temperature for 2 h as described above in EXAMPLES 7-8. The product
was collected by filtration from the aqueous work-up. This material was
dissolved in isopropanol (~30 mL) and treated with a saturated solution of
hydrochloric acid in isopropanol (~ 5 mL). The solvent was evaporated *in vacuo*, and the residue was triturated with diethyl ether to give 4-acetyl-4-phenyl-N-[cis-1,2,3,4-tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-
15 piperidineacetamide bis-hydrochloride **21** as an amorphous pale yellow solid
(1.2 g, 90%): MS *m/e* 482 (MH⁺) (Figure 7).




Figure 7

EXAMPLE 10

4-Oxo-1-phenyl-*N*-[*cis*-1,2,3,4-tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenyl]-
5 1,3,8-triazaspiro[4.5]decane-8-acetamide bis-hydrochloride 22

1,2,3,4-Tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenamine bis-hydrochloride
4 (0.75 g, 2.41 mmol) was reacted with 2-(1-phenyl-1,3,8-triaza-
spiro[4.5]decan-4-one)acetic acid (1.12 g, 2.41 mmol), *N,N*-
10 diisopropylethylamine (1.68 mL, 9.63 mmol) and HBTU (0.91 g, 2.41
mmol) in *N,N*-dimethylformamide (15 mL) at room temperature for 4 h as
described above in EXAMPLES 7-8. The product was collected by filtration
from the aqueous work up. This material was dissolved in methanol (~30 mL),
and treated with concentrated hydrochloric acid (~ 5 mL). The solvent was
15 evaporated *in vacuo*, and the residue was triturated with diethyl ether to give 4-
oxo-1-phenyl-*N*-[*cis*-1,2,3,4-tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenyl]-
1,3,8-triazaspiro[4.5]decane-8-acetamide bis-hydrochloride 22 as an
amorphous tan solid (1 g, 81%): ¹H NMR(DMSO-*d*₆) δ 1.93 (s, 4 H), 2.80-3.08
(m, 4 H), 3.18-3.30 (m, 2 H), 3.38-3.66 (m, 3 H), 3.70-3.89 (m, 2 H), 3.94-4.13
20 (m, 3 H), 4.65 (s, 2 H), 6.80 (t, 2 H), 7.00-7.29 (m, 8 H), 8.03 (t, 1 H), 8.44 (d, 1
H), 8.81 (br s, 2 H), 8.97 (d, 1 H), 9.16 (s, 1 H), 10.83 (br s, 1 H); MS *m/e* 510
(MH⁺) (Figure 8).




Figure 8

5 EXAMPLE 11

4-(2,3-Dihydro-2-oxo-1H-benzimidazol-1-yl)-N-[cis-1,2,3,4-tetrahydro-6-hydroxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-hydrochloride 23

10

A solution of 4-(2,3-dihydro-2-oxo-1H-benzimidazol-1-yl)-N-[cis-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide 19 (0.28 g, 0.37 mmol) in dichloromethane (2 mL) was added dropwise to a solution of boron tribromide (1.8 mmol) in dichloromethane (22 mL) at 0°C. After stirring the resultant solution at 0°C for 1.5 h, methanol (~2 mL) was added and stirring was continued at 0°C for an additional 0.5 h. The solvent was evaporated *in vacuo*, and the residue was purified by reverse phase C₁₈ HPLC using a water/acetonitrile/TFA gradient, 90/10/0.1 to 10/90/0.1, as the eluant. The product was dissolved in methanol

15

and treated with ethanolic hydrochloric acid. The solvent was evaporated and the process repeated twice to give 4-(2,3-dihydro-2-oxo-1*H*-benzimidazol-1-yl)-*N*-[*cis*-1,2,3,4-tetrahydro-6-hydroxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-hydrochloride salt **23** (0.148, 68%) as a colorless solid:

5 ¹H NMR(DMSO-*d*₆) δ 1.73-2.03 (m, 4 H), 2.70-2.94 (m, 4 H), 3.05-3.20 (br s, 2 H), 3.27-3.47 (m, 3 H), 3.55-3.76 (m, 2 H), 3.92-4.15 (m, 3 H), 4.54-4.67 (m, 1 H), 6.46 (d, 1 H), 6.58 (s, 2 H), 7.05 (m, s, 3 H), 7.60 (br s, 1 H), 7.94 (t, 1 H), 8.30 (d, 1 H), 8.72-8.83 (m, 2 H), 8.96 (d, 1 H), 9.30 (br s, 1 H), 10.64 (br s, 1 H), and 11.05 (s, 1 H); MS *m/e* 512 (MH⁺) (Figure 9).

10




Figure 9

15

EXAMPLES 12-13

trans-N-[2-(4-fluorophenyl)-3-(3-pyridinyl)propyl]-4-[[2-fluorophenylsulfonyl)amino)methyl]-1-cyclohexanamide hydrochloride **26**

5

trans-N-[[[2-(4-fluorophenyl)-3-(3-pyridinyl)propyl]amino)methyl]-4-cyclohexyl)methyl] 2-fluorobenzenesulfonamide bis-hydrochloride **27**

A. Sodium metal (0.71 g, 30.9 mmol) was added to methanol (75 mL) and stirred at room temperature until the solid was consumed. At this time, 4-fluorophenylacetonitrile (3.5 mL, 29.3 mmol) was added and the mixture was stirred at room temperature for 10 min. 3-Pyridinecarboxaldehyde (2.77 mL, 29.3 mmol) was added and the resultant solution was heated at reflux for 2 h. The reaction was cooled to room temperature and neutralized with 2 N hydrochloric acid (16 mL, 32 mmol). The solvent was evaporated *in vacuo*, and the resultant residue was partitioned between water (~200 mL) and dichloromethane (~200 mL). The organic layer was dried over sodium sulfate, filtered and the solvent was evaporated *in vacuo* to give 2-(4-fluorophenyl)-3-pyridin-3-yl-acrylonitrile **24** as a colorless solid (6.11 g, 93%); ¹H NMR(CDCl₃) δ 7.16 (t, 2 H), 7.42-7.47 (m, 1 H), 7.48 (s, 1 H), 7.66-7.70 (m, 2 H), 8.47 (d, 1 H), 8.65 (d, 1 H), 8.84 (s, 1 H); MS *m/e* 225 (MH⁺)

B. A suspension of 2-(4-fluorophenyl)-3-pyridinyl-3-acrylonitrile **24** (1.5 g, 6.68 mmol) and platinum(IV) oxide (0.51 g, 2.24 mmol) in ethanol (60 mL) and water (15 mL) was reacted with hydrogen gas at a pressure of 65 psi for 6 h. The catalyst was removed by filtration, and the solvent was evaporated *in vacuo*. The residue was dissolved in diethyl ether (50 mL), and the small amount of insoluble material was removed by filtration. The ethereal solution was treated with 1 N hydrogen chloride in diethyl ether (20 mL). A yellow solid precipitated which was collected by filtration and washed generously with diethyl ether to give β-(3-pyridinylmethyl)-4-fluorophenethylamine bis hydrochloride salt **25** as a pale yellow solid (1.67 g, 82%). ¹H NMR(DMSO-d₆)

δ 3.03-3.21 (m, 4 H), 3.44-3.53(m, 1 H), 7.13 (t, 2 H), 7.27-7.33 (m, 2 H), 7.93 (t, 1 H), 8.27 (d, 1 H), 8.42 (br s, 3 H), 8.72-8.80 (m, 2 H); MS m/e 231 (MH^+).

- 5 C. A solution of 2-(1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) (1.03 g, 2.57 mmol), *trans*-4-[(2-fluorophenyl)sulfonylamino]methyl]cyclohexanecarboxylic acid (1.20 g, 2.57 mmol), and *N,N*-diisopropylethylamine (1.9 mL, 11.1 mmol) in *N,N*-dimethylformamide (15 mL) was stirred at room temperature for 10 min. 2-(4-
10 Fluorophenyl)-3-pyridin-3-yl-propylamine dihydrochloride **25** (0.75 g, 2.47 mmol) was added, and the resultant solution was stirred at room temperature for 2 h. The reaction mixture was poured into water (~100 mL) and the product was extracted into dichloromethane (~100 mL). The organic layer was washed with water (3 x 100 mL), concentrated and the resultant residue purified via
15 flash chromatography using methanol (5-10%) and triethylamine (0.5%) in dichloromethane as the eluant to give the desired cyclohexanamide as an oil. This material was dissolved in diethyl ether (~50 mL) and treated with 1 N hydrogen chloride in diethyl ether. A colorless solid formed which was collected by filtration, washed with ether and dried *in vacuo* to give *N*-[2-(4-
20 fluorophenyl)-3-(3-pyridinyl)propyl]-4-[(2-fluorophenylsulfonyl)amino]methyl]-1-cyclohexanamide hydrochloride **26** as a colorless solid. 1H NMR(DMSO- d_6) δ 0.69-0.83 (m, 2 H), 1.07-1.19 (m, 3 H), 1.52-1.71 (m, 4 H), 1.94 (t, 1 H), 2.66 (br s, 2 H), 2.99-3.10 (m, 1 H), 3.17-3.43 (m, 4 H), 7.07 (t, 2 H), 7.16-7.21 (m, 2 H), 7.35-7.47 (m, 2 H), 7.66-7.95 (m, 5 H), 8.28 (d, 1 H), and 8.74 (br s, 2 H);
25 MS m/e 528 (MH^+) (Figure 10).

- D. *N*-[2-(4-Fluorophenyl)-3-(3-pyridinyl)propyl]-4-[(2-fluorophenylsulfonyl)amino]methyl]-1-cyclohexanamide hydrochloride **26** was
30 partitioned between a saturated solution of aqueous sodium bicarbonate and dichloromethane. The organic layer was dried over sodium sulfate and the solvent was evaporated *in vacuo* to give the free base as an oil. This oil (0.5 g, 0.944 mmol) was dissolved in tetrahydrofuran (~20 mL), and the resultant

solution was added dropwise to a solution of borane (4.0 mmol) in tetrahydrofuran (14 mL) at ambient temperature. The solution was heated at reflux for 2 h. The resultant mixture was cooled to room temperature and several drops of water were added until unreacted borane was consumed. A
5 4 N solution of hydrochloric acid (2 mL) was added and the solution heated at reflux for 45 min. After the solution had cooled, 3 N aqueous sodium hydroxide was added (2.7 mL), and the mixture was concentrated *in vacuo*. The residue was partitioned between water (~50 mL) and dichloromethane (~50 mL). The organic layer was dried over sodium sulfate, and the solvent was evaporated *in*
10 *vacuo*. The residue was dissolved in diethyl ether (~20 mL) and treated with 1 N hydrogen chloride in diethyl ether (~4 mL). The colorless precipitate was collected by filtration, washed generously with diethyl ether and dried *in vacuo* to give *trans-N*-[[[2-(4-fluorophenyl)-3-(3-pyridinyl)propyl]amino]methyl]-4-cyclohexyl]methyl] 2-fluorobenzenesulfonamide bis-hydrochloride **27** (0.371 g,
15 67%): ¹H NMR(DMSO-d₆) δ 0.70-0.87 (m, 4 H), 1.22-1.36 (br s, 1 H), 1.64-1.88 (m, 6 H), 2.65-2.77 (m, 3 H), 2.99-3.33 (m, 3 H), 3.54-3.70 (m, 2 H), 7.13 (t, 2 H), 7.24 -7.34 (m, 2 H), 7.37-7.48 (m, 2 H), 7.67-7.87 (m, 3 H), 7.96 (t, 1 H), 8.17 (d, 1 H), 8.68 (s, 1 H), 8.70 (s, 1 H), 9.03 (br s, 1 H), and 9.24 (br s, 1 H); MS *m/e* 514 (MH⁺) (Figure 10).

20




Figure 10

5 EXAMPLE 14

N-[2-(4-Fluorophenyl)-3-(3-pyridinyl)propyl]-4-[(2-fluorophenylsulfonyl)amino]-1-piperidineacetamide bis-trifluoroacetate **30**

- 10 A. A solution of [4-(1,1-dimethylethoxy)carbonylamino-piperidin-1-yl]acetic acid (0.5 g, 1.94 mmol), 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (0.73 g, 1.94 mmol), and *N,N*-diisopropylethylamine (1.5 mL, 8.71 mmol) in *N,N*-dimethylformamide (15 mL) was stirred at room temperature for 5 min. β -(3-Pyridinylmethyl)-4-fluorophenethylamine
- 15 dihydrochloride **25** (0.586 g, 1.94 mmol) was added, and the resultant solution

- was stirred at room temperature for 24 h. The solution was poured into a saturated solution of aqueous sodium bicarbonate (~100 mL) and the product was extracted into dichloromethane (~100 mL). The organic layer was washed with water (5 x ~100 mL) and dried over sodium sulfate. The solvent was
- 5 evaporated *in vacuo* to give the piperidineacetamide **28** as an oil, 0.52 g (57%): ¹H NMR(CDCl₃) δ 0.98-1.25 (m, 2 H), 1.45 (s, 9 H), 1.71-1.79 (m, 2 H), 2.05-2.17 (m, 2 H), 2.41-2.50 (m, 2 H), 2.75-3.00 (m, 3 H), 3.04-3.17 (m, 1 H), 3.33-3.47 (m, 2 H), 3.72-3.83 (m, 1 H), 4.36 (br s, 1 H), 6.93-7.14 (m, 7 H), 7.25 (m, 1 H), 8.24 (s, 1 H), 8.39 (d, 1 H); MS *m/e* 471 (MH⁺).
- 10
- B. A solution of the piperidineacetamide **28** (0.46 g, 0.977 mmol) in dichloromethane (6 mL) was treated with trifluoroacetic acid (2 mL) and stirred at room temperature for 3 h. The solvent was evaporated *in vacuo*. The residue was dissolved in 1,2-dichloroethane (~10 mL), and the solvent
- 15 evaporated *in vacuo* (repeated twice to remove residual trifluoroacetic acid), to give the 4-amino-1-piperidineacetamide **29** as a tris-trifluoroacetate salt, isolated as an amber glass, 0.66 g (95%): ¹H NMR(DMSO-d₆); MS *m/e* 371 (MH⁺).
- 20
- C. 2-Fluorobenzenesulfonyl chloride (25 mg, .126 mmol) was added to a solution of the 4-amino-1-piperidineacetamide **29** (82 mg, 0.115 mmol) and *N,N*-diisopropylethylamine (0.10 mL, 0.575 mmol) in acetonitrile (1 mL) at room temperature. The mixture was stirred at room temperature for 16 h and then water (0.30 mL) was added and the solution was applied to a C₁₈ reverse
- 25 phase column for purification by HPLC. The column was eluted with a gradient of water/acetonitrile/trifluoroacetic acid to give N-[2-(4-fluorophenyl)-3-(3-pyridinyl)propyl]-4-[(2-fluorophenylsulfonyl)amino]-1-piperidineacetamide bis-trifluoroacetate **30** as a colorless solid, 28 mg (32%): ¹H NMR(DMSO-d₆) δ 1.70-1.85 (m, 4 H), 2.91-3.47 (m, 10 H), 3.66-3.80 (m, 2 H), 7.07 (t, 2 H), 7.18 (m, 2 H), 7.38-7.50 (m, 2 H), 7.64 (t, 1 H), 7.71-7.85 (m, 2 H), 7.92 (d, 1 H),
- 30 8.31 (d, 1 H), 8.49 (s, 1 H), 8.57 (s, 1 H), 8.60 (s, 1 H); MS *m/e* 529 (MH⁺) (Figure 11).





Figure 11

5

Additional compounds of this invention that were prepared using the experimental protocols described above include:



























Mass Spectral Data of Compounds





























A































5































#	R ₁	R ₂	m	B ₁	B ₂	Y	L	Z	MH +	Calc M
31	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			496	495
19	6-OMe	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			526	525
20	6-OMe	3-pyridyl (trans)	1	-CH ₂ -	-CH ₂ -	C=O			526	525
23	6-OH	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			512	511
32	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			525	524
33	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			507	506
34a	(H)	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	C=O			507	506
34b	(H)	3-pyridyl (diast-B)	1	-CH ₂ -	-CH ₂ -	C=O			507	506

























35	6-OMe	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			555	554
7a	6-OMe	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	C=O			555	554
7b	6-OMe	3-pyridyl (diast-B)	1	-CH ₂ -	-CH ₂ -	C=O			555	554
8a	6-OMe	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			541	540
9a	6-OH	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			527	526
36	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			567	566
37	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			549	548
38a	(H)	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	C=O			549	548
38b	(H)	3-pyridyl (diast-B)	1	-CH ₂ -	-CH ₂ -	C=O			549	548
10	6-OMe	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			597	596
39	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			611	610
40	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			578	577
41	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			550	549



















WO 01/09120

42	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			549	548
43a	(H)	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	C=O			535	534
43b	(H)	3-pyridyl (diast-B)	1	-CH ₂ -	-CH ₂ -	C=O			535	534
11a	6-OH	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	C=O			583	582
11b	6-OH	3-pyridyl (diast-B)	1	-CH ₂ -	-CH ₂ -	C=O			583	582
17	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			523	522
44	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			501	500
45	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			516	515
46	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			517	516
47	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			515	514
48	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			537	536
49	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			555	554
22	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			510	509

21	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			482	481
50	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			537	536
51	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			498	497
52	6-OMe	3-thienyl	1	-CH ₂ -	-CH ₂ -	C=O			572	571
53	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			496	495
54	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			497	496
55	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			483	482
56	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			483	482
57	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			483	482
58	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			523	522
59	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			523	522
60	6-F	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			487	486
61	6-OMe	3-thienyl	1	-CH ₂ -	-CH ₂ -	C=O			531	530
62	6-OMe	3-thienyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			517	516
63	6-OMe	3-thienyl	1	-CH ₂ -	-CH ₂ -	C=O			560	559

64	6-OMe	3-thienyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			546	545
65	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			493	492
66	(H)	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			493	492
67	6-F	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			529	528
68	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			521	520
69	6-OMe	5(4)- imidazol yl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			530	529
70	6-F	3-pyridyl (diast-A)	1	-CH ₂ -	-CH ₂ -	C=O			543	542
71	6-F	3-pyridyl (diast-B)	1	-CH ₂ -	-CH ₂ -	C=O			543	542
72	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			626	625
73	6-OMe	5(4)- imidazol yl	1	-CH ₂ -	-CH ₂ -	C=O			515	514
74	6-OMe	3-thienyl	1	-CH ₂ -	-CH ₂ -	C=O			602	601
75	6-OMe	4-Cl- phenyl	1	-CH ₂ -	-CH ₂ -	C=O			588	587
76	6-OMe	4-Cl- phenyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			574	573
77	6-F	vinyl	1	-CH ₂ -	-CH ₂ -	C=O			474	473
78	6-F	vinyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			460	459

79	6-OMe	vinyl	1	-CH ₂ -	-CH ₂ -	C=O			475	474
80	6-OMe	vinyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			461	460
81	6-OH	vinyl	1	-CH ₂ -	-CH ₂ -	-CH ₂ -			447	446
82	6-OMe	(H)	0	-CH ₂ -	-CH ₂ -	C=O			435	434
83	6-OH	(H)	0	-CH ₂ -	-CH ₂ -	C=O			421	420
84	6-OMe	(H)	0	-CH ₂ -	-CH ₂ -	-CH ₂ -			461	460
85	6-OH	(H)	0	-CH ₂ -	-CH ₂ -	-CH ₂ -			447	446
86	6-OMe	3-pyridyl	1	H	H	C=O			500	499
87	6-OH	3-pyridyl	1	H	H	C=O			486	485
88	6-OMe	3-pyridyl	1	H	H	C=O			540	539
89	6-OMe	3-pyridyl	1	H	H	-CH ₂ -			526	525
90	6-OH	3-pyridyl	1	H	H	C=O			526	525

91	6-OH	3-pyridyl	1	H	H	-CH ₂ -			512	511
26	6-F	3-pyridyl	1	H	H	C=O			528	527
27	6-F	3-pyridyl	1	H	H	-CH ₂ -			514	513
92	6-F	3-pyridyl	1	H	H	C=O			475	474
30	6-F	3-pyridyl	1	H	H	C=O			529	528
93	(H)	3-pyridyl	1	-CH ₂ -	-CH ₂ -	C=O			471	470
94	6-OMe	(H)	0	H	H	C=O			449	448
95	6-OMe	(H)	0	H	H	-CH ₂ -			435	434
96	6-OH	(H)	0	H	H	-CH ₂ -			421	420

IN VITRO ASSAYS

5

NPY5 HTS Centrifugation Assay

The compounds described in this invention were evaluated for binding to the human neuropeptide Y5 receptor.

10

Stable Transfection

The human NPY5 receptor cDNA (Genbank Accession number U66275) was inserted into the vector pCIneo (Invitrogen) and transfected into

human embryonic kidney cells (HEK-293) via Calcium phosphate method (Cullen 1987). Stably transfected cells were selected with G-418 (600 ug/mL). Stably transfected cells served as the source for the membranes for the NPY5 receptor binding assay.

5

Membrane Preparation

NPY5-transfected HEK293 cells were grown to confluence in 150 cm² culture dishes. Cells were washed once with phosphate-buffered saline (Gibco Cat# 14040-133). Cells were then incubated in phosphate-buffered saline without Calcium and without Magnesium, supplemented with 2 mM EDTA. Cells were incubated for 10 minutes at room temperature and the cells were collected by repetitive pipeting. Cells were formed into pellets and then frozen at -80 until needed. Frozen pellets were homogenized with a polytron at full speed for 12 seconds in a homogenization buffer (20 mM Tris HCl, 5 mM EDTA, pH 7.4). Homogenates were centrifuged for 5 minutes at 4C at 200g. Supernatants were transferred to corex tubes and centrifuged for 25 minutes at 28,000g. Pellets were re-suspended in Binding (20mM HEPES, 10 mM NaCl, 0.22 mM KH₂PO₄, 1.3mM CaCl₂, 0.8 mM MgSO₄, pH 7.4). Membranes were kept on ice until use.

A competition binding assay, known to those skilled in the art, was used in which compounds of formula A compete with ¹²⁵I-PYY for binding to cell membranes. In simple terms, the less ¹²⁵I-PYY bound to the membranes implies that a compound is a good inhibitor (competitor). Bound ¹²⁵I-PYY is determined by centrifugation of membranes, aspirating supernatant, washing away residual ¹²⁵I-PYY and subsequently counting the bound sample in a g-counter.

30 Procedure for Radioligand binding assay


Compounds to be tested were prepared as 10x stocks in binding buffer and added first to assay tubes (RIA vials, Sarstedt). Twenty (20) µL of each

WO 01/09120

10x compound stock is pipeted into vials and 80 μ L of 125 I-PYY (NEN catalog number NEX240), which has been diluted to a concentration of 200 pM in 0.25 % BSA in binding buffer, is added to the compound tubes (final concentration of 125 I-PYY is 80 pM). To each tube is added 100 μ L of membranes and the mixture is agitated by pipeting 2 times. Samples are incubated for 1 hr at room temperature. Aluminum cast plates (Sarstedt) containing the vials are then centrifuged 10 minutes at 3200 rpm in a Sorvall RT6000. Supernatant is then aspirated. To each vial 400 μ L PBS is added and this is then aspirated again. Vials are then put in carrier polypropylene 12x75 tube and counted in gamma counter (Packard). Non-specific binding is determined in the presence of 300 nM NPY. Percent inhibition of 125 I-PYY binding is calculated by subtracting non-specific binding from the test samples (compound (I)), taking these counts and dividing by total binding, and multiplying by 100. Inhibitory concentration values (IC_{50}) of compounds that show appreciable inhibition of 125 I-PYY binding are calculated by obtaining percent inhibition of 125 I-PYY binding values at different concentrations of the test compound, and using a graphing program such as GraphPad Prism (San Diego, CA) to calculate the concentration of test compound that inhibits fifty-percent of 125 I-PYY binding (Table 4). These operations are known to those skilled in the art.

20

Binding Affinities of Compounds A for the Human NPY Y5 Receptor
 (expressed as % Inhibition of ¹²⁵I-PYY Binding)



A

5

#	%Inh @ 3 uM	%Inh @ 300 nM
7a	97	69
7b	67	11
8	100	96
9	98	104
10	96	60
17	102	98
19	101	69
20	96	88
21	98	83
22	70	32
23	100	96
26	110	108
27	110	105
30	110	100
31	100	91
32	100	62
33	96	52
34a	97	87
34b	99	61
35	96	54
36	95	22
37	102	89
38a	104	80
38b	101	89
39	95	70

40	92	21
41	94	54
42	85	21
43a	93	84
43b	86	62
44	98	93
45	95	68
46	107	90
47	98	91
48	103	97
49	95	85
50	108	103
51	102	85
52	100	96
53	92	84
54	100	99
55	106	96
56	94	88
57	93	87
58	91	93
59	93	90
60	109	86
61	87	66
62	103	74
63	71	33
64	103	91
65	98	79
66	102	98
67	99	102
68	108	109
69	56	26
70	92	93
71	73	59
72	73	41
73	63	32
74	100	89
75	78	28
76	91	45

77	84	56
78	75	65
79	99	69
80	82	47
81	94	89
82	85	63
83	92	72
84	93	79
85	100	96
86	91	88
87	96	97
88	103	104
89	100	103
90	88	93
91	100	104
92	104	92
93	97	81
94	98	93
95	102	96
96	98	91

Table 2

IN VIVO ASSAYS

5

Rodent Feeding Model:

Measurement of Food Intake in Food-Deprived Rats

Male Long-Evans rats (180-200 grams) are housed individually and are maintained on a once-a-day feeding schedule (i.e. 10 a.m. until 4 p.m.) for five days following quarantine to allow the animals to acclimate to feeding on powdered chow (#5002 PMI Certified Rodent Meal) during the allotted time. The chow is made available in an open jar, anchored in the cage by a wire, with a metal follower covering the food to minimize spillage. Water is available ad-libitum.

15

WO 01/09120

Animals are fasted for 18 hours prior to testing. At the end of the fasting period, animals are administered either compounds of the invention or vehicle. Vehicle and test compounds are administered either orally (5 mL/kg) 60 minutes prior to the experiment, or 30 minutes prior when given subcutaneously (1 mL/kg) or intraperitoneally (1 mL/kg). Compounds of the invention are administered orally as a suspension in aqueous 0.5% methylcellulose-0.4% Tween 80, or intraperitoneally as a solution or suspension in PEG 200; compound concentrations typically range from 1 mg/kg to 100 mg/kg, preferably from 10-30 mg/kg. Food intake is measured at 2, 4, and 6 hours after administration by weighing the special jar containing the food before the experiment and at the specified times. Upon completion of the experiment, all animals are given a one-week washout period before retesting.

Percent reduction of food consumption is calculated subtracting the grams of food consumed by the treated group from the grams of food consumed by the control group divided by the grams of food consumed by the control group, multiplied by 100.

$$\% \text{ change} = \frac{\text{Treatment} - \text{Vehicle}}{\text{Vehicle}} \times 100$$


A negative value indicates a reduction in food consumption and a positive value indicates an increase in food consumption.

Compound	Dose (mg/kg)	Food Consumption (grams)			
		2 hrs	4 hrs	6 hrs	2-6 hrs
	(#rats)	(%chg.)	(%chg.)	(%chg.)	(%chg.)
Vehicle	N=6	8.85 g	13.97 g	22.85 g	14.00 g
PEG-2000					
70	30 (i.p.)	1.30 g	3.44 g	6.14 g	4.84 g
	N=7	(-85.3%)	(-75.4%)	(-73.1%)	(-65.4%)

What is claimed is:

1. A compound of the formula:

5



A

10 in which:

R_1 is independently selected from the group consisting of hydrogen; hydroxy; halo; C_{1-6} alkyl; substituted C_{1-6} alkyl; C_{1-8} alkoxy; substituted C_{1-8} alkoxy; trifluoroalkyl; C_{1-8} alkylthio; substituted C_{1-8} alkylthio; C_{3-6} cycloalkyl; C_{3-8} cycloalkoxy; nitro; amino; C_{1-6} alkylamino; C_{1-8} dialkylamino; C_{4-8} cycloalkylamino; cyano; carboxy; C_{1-5} alkoxycarbonyl; C_{1-5} alkylcarbonyloxy; formyl; carbamoyl; phenyl; and substituted phenyl;

20 n is 1-2;

B_1 is hydrogen;

25 B_2 is hydrogen; or B_1 and B_2 are methylene and joined together to form a five or six membered ring;


m is 0-3;

R_2 is independently selected from the group consisting of hydrogen; hydroxy; C_{1-6} alkyl; C_{2-6} alkenyl; halo; C_{3-7} cycloalkyl; phenyl; substituted phenyl; naphthyl, substituted naphthyl; phenoxy; substituted phenoxy; heteroaryl; substituted heteroaryl; and heterocycloalkyl;

5


L is selected from the group consisting of C_{1-8} alkylene; C_{2-10} alkenylene; C_{2-10} alkynylene; C_{3-7} cycloalkylene; C_{3-7} cycloalkyl C_{1-4} alkylene; aryl C_{1-4} alkylene; α -amino C_{4-7} alkylene;

10




(N-methylene)piperidin-4-yl;

15




(N-methylene)piperazin-4-yl;

20




(N-methylene)pyrrolidin-3-yl;




(N-methylene)-4-acetyl-piperidin-4-yl;

25



and (N-methylene)piperidin-4,4-diyl;




5 Y is methylene or carbonyl;


Z is selected from the group consisting of:

aryl;

10




N-sulfonamido;




15

N-(aryl)sulfonamido;




arylamido;




20


arylureido;



arylacetamido:




(aryloxy)carbonylamino;




5

2,3-dihydro-2-oxo-1*H*-benzimidazol-1-yl;



and 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl;

10



15 R_3 is independently selected from the group consisting of C_{1-8} alkyl; substituted C_{1-8} alkyl; cycloalkyl; substituted cycloalkyl; naphthyl; substituted naphthyl; heteroaryl; and substituted heteroaryl;

20 R_4 is independently selected from the group consisting of hydrogen; C_{1-8} alkyl; C_{1-8} alkoxy; substituted C_{1-8} alkoxy; hydroxy; halogen; cyano; nitro; amino; C_{1-8} alkylamino; and C_{1-8} dialkylamino;

R_5 is independently selected from the group consisting of hydrogen;
 C_{1-8} alkyl; C_{1-8} alkylcarbonyl; aroyl; carbamoyl; amidino; C_{1-8} alkyl;
 C_{1-8} alkylaminocarbonyl; (arylamino)carbonyl; and aryl C_{1-8} alkylcarbonyl;

5 R_6 is independently selected from hydrogen and C_{1-8} alkyl;

p is 1-3;

q is 1-3;

10

and enantiomers, diastereomers and pharmaceutically acceptable salts thereof;

provided that when L is C_{1-6} alkylene; C_{2-10} alkenylene; C_{2-10} alkynylene;

15 C_{3-7} cycloalkylene; C_{3-7} cycloalkyl C_{1-4} alkylene; aryl C_{1-4} alkylene; or α -amino C_{4-7} alkylene; then Z is phenyl, N-sulfonamido or N-(aryl)sulfonamido;

when L is (N-methylene)piperazin-4-yl; then Z is phenyl or naphthyl;

20 when L is (N-methylene)pyrrolidin-3-yl or (N-methylene)piperidin-4-yl; then Z is N-sulfonamido; N-(aryl)sulfonamido; 2,3-dihydro-2-oxo-1*H*-benzimidazol-1-yl; benzamido; phenylureido; phenylacetamido or (phenoxy)carbonylamino;

when L is (N-methylene)-4-acetyl-piperidin-4-yl; then Z is phenyl or naphthyl

25 and Y is carbonyl;

when L is (N-methylene)piperidin-4,4-diyl; then Z is 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl and Y is carbonyl;


30 and when B_1 and B_2 are both methylene thus forming a six membered ring and when L is C_{1-10} alkylene; C_{2-10} alkenylene; C_{2-10} alkenylene; or aryl C_{1-4} alkylene; then Z is other than N-sulfonamido, N-(aryl)sulfonamido or phenyl.

2. A compound of claim 1 wherein R_1 is H; alkyl; substituted alkyl; alkoxy; halo; substituted alkoxy; hydroxy; trifluoralkyl; nitro; amino; alkylamino; cycloalkylamino; cyano; carboxy; cycloalkyl; phenyl; and substituted phenyl;
- 5 R_2 is H; hydroxy; alkyl; substituted alkyl; halo; heterocycloalkyl; heteroaryl; phenyl; substituted phenyl; naphthyl and substituted naphthyl;
- B_1 is hydrogen;
- 10 B_2 is hydrogen; or B_1 and B_2 are methylene and joined together to form a five or six membered ring;
- Y is methylene or carbonyl;
- 15 Z is selected from the group consisting of aryl; substituted aryl; N-sulfonamido; N-(aryl)sulfonamido; substituted N-(aryl)sulfonamido; arylamido; substituted arylamido; arylureido; substituted arylureido; arylacetamido; substituted arylacetamido; (aryloxy)carbonylamino; substituted (aryloxy)carbonylamino;
- 20 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl; substituted 2,3-dihydro-2-oxo-1H-benzimidazol-1-yl; 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl; and substituted 1-aryl-2,3-dihydro-4-oxo-imidazol-5,5-diyl;
- L is C_{1-6} alkylene; C_{2-10} alkenylene; C_{2-10} alkynylene; C_{3-7} cycloalkylene;
- 25 C_{3-7} cycloalkyl C_{1-4} alkylene; aryl C_{1-4} alkylene; α -amino C_{4-7} alkylene; (N-methylene)piperidin-4-yl; substituted (N-methylene)piperidin-4-yl; (N-methylene)piperazin-4-yl; (N-methylene)pyrrolidin-3-yl; (N-methylene)-4-acetyl-piperidin-4-yl; and (N-methylene)piperidin-4,4-diyl;
- 30 n is 1-2;
- m is 1-3;







p is 1-3; and

q is 1-3.

5 3. A compound of claim 1 selected from the group consisting of:









4. A compound of claim 1 selected from the group consisting of:



and


5. A compound of claim 1 selected from the group consisting of:



and


WO 01/09120

6. A compound of claim 1 selected from the group consisting of:








and

7. A compound of claim 1 selected from the group consisting of:








and

8. A compound of claim 1 selected from the group consisting of:








and

9. A compound of claim 1 selected from the group consisting of:




and







10. A compound of claim 1 selected from the group consisting of:



and




11. A compound of claim 1 selected from the group consisting of:



and

12. A compound of claim 1 selected from the group consisting of:



and

13. A compound of claim 1 selected from the group consisting of:

2-Amino-6-[(2-fluorophenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthenyl-(2*S*)-hexanamide bis-hydrochloride,

N-[5-amino-6-[[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenyl]amino]hexyl-2-fluorobenzenesulfonamide tris-hydrochloride,

N-[5-amino-6-[[*cis*-1,2,3,4-tetrahydro-6-hydroxy-1-(3-pyridinylmethyl)-2-naphthalenyl]amino]hexyl-2-fluorobenzenesulfonamide tris-hydrochloride,

(2*S*)-2-(Acetylamino)-6-[(2-fluorophenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthenyl]hexanamide bis-hydrochloride,

(2*S*)-2-(Acetylamino)-6-[(2-fluorophenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-hydroxy-1-(3-pyridinylmethyl)-2-naphthenyl]hexanamide bis-hydrochloride,

3-[(Phenylsulfonyl)amino]-N-[*cis*-1,2,3,4-tetrahydro-6-fluoro-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-pyrrolidineacetamide bis-trifluoroacetate,

4-(2,3-Dihydro-2-oxo-1*H*-benzimidazol-1-yl)-N-[*cis*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-hydrochloride,

4-(2,3-Dihydro-2-oxo-1*H*-benzimidazol-1-yl)-N-[*trans*-1,2,3,4-tetrahydro-6-methoxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-hydrochloride,

4-Acetyl-4-phenyl-N-[*cis*-1,2,3,4-tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-hydrochloride,

5 4-Oxo-1-phenyl-N-[*cis*-1,2,3,4-tetrahydro-1-(3-pyridinylmethyl)-2-naphthalenyl]-1,3,8-triazaspiro[4.5]decane-8-acetamide bis-hydrochloride,

10 4-(2,3-Dihydro-2-oxo-1*H*-benzimidazol-1-yl)-N-[*cis*-1,2,3,4-tetrahydro-6-hydroxy-1-(3-pyridinylmethyl)-2-naphthalenyl]-1-piperidineacetamide bis-hydrochloride,

trans-N-[2-(4-fluorophenyl)-3-(3-pyridinyl)propyl]-4-[(2-fluorophenylsulfonyl)amino)methyl]-1-cyclohexanamide hydrochloride,

15 *trans*-N-[[[2-(4-fluorophenyl)-3-(3-pyridinyl)propyl]amino)methyl]-4-cyclohexyl]methyl] 2-fluorobenzenesulfonamide bis-hydrochloride and

20 N-[2-(4-fluorophenyl)-3-(3-pyridinyl)propyl]-4-[(2-fluorophenylsulfonyl)amino]-1-piperidineacetamide bis-trifluoroacetate.

14. A method of treating disorders and diseases associated with NPY receptor subtype 5 comprising administering to a mammal in need of such treatment a therapeutically effective amount of a compound of claim 1.

25 15. A pharmaceutical composition for the treatment of diseases or disorders associated with the NPY Y5 receptor subtype comprising a therapeutically effective amount of a compound of claim 1 and a pharmaceutically acceptable carrier.

30 16. A pharmaceutical composition according to claim 15 for the treatment of disorders or disease states caused by eating disorders, obesity, bulimia

nervosa, diabetes, memory loss, epileptic seizures, migraine, sleep disturbances, pain, sexual/reproductive disorders, depression and anxiety.

INTERNATIONAL SEARCH REPORT

Intern I Application No
PCT/US 00/20482

A. CLASSIFICATION OF SUBJECT MATTER			
IPC 7	C07D401/12	C07D401/14	C07D213/40
	C07D401/04	C07D211/34	C07D295/14
	C07D333/20	A61K31/444	A61K31/4523
			A61P25/00
			C07D471/10
			C07D213/38
			C07C311/18
			C07D409/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category * Citation of document, with indication, where appropriate, of the relevant passages

Relevant to claim No.

X,P WO 99 55667 A (ORTHO MCNEIL PHARM INC)
4 November 1999 (1999-11-04)
the whole document

A WO 98 35957 A (BAYER AG)
20 August 1998 (1998-08-20)
cited in the application
claims; examples

A WO 97 19682 A (FINN JOHN M ;JEON YOON T
(US); PING DU (US); DHANOA DALJIT S (US);)
5 June 1997 (1997-06-05)
cited in the application
claims; examples

1,14-16

1,14-16

1,14-16

-/--

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

17 November 2000

Date of mailing of the international search report

05/12/2000

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bosma, P

INTERNATIONAL SEARCH REPORT

Inte... Application No

PCT/US 00/20482

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	<p>M.A. YOUNGMAN ET AL.: "Alpha-substituted N-(sulfonamido)alkyl-beta-aminotetralins: potent and selective neuropeptide Y Y5 receptor antagonists" JOURNAL OF MEDICINAL CHEMISTRY., vol. 43, no. 3, February 2000 (2000-02), pages 346-350, XP002153193 AMERICAN CHEMICAL SOCIETY. WASHINGTON., US ISSN: 0022-2623 the whole document</p>	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/20482

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9955667 A	04-11-1999	AU 3640099 A US 6140354 A	16-11-1999 31-10-2000
WO 9835957 A	20-08-1998	AU 6144098 A CA 2251368 A EP 0910565 A US 5939462 A	08-09-1998 20-08-1998 28-04-1999 17-08-1999
WO 9719682 A	05-06-1997	AU 1328197 A	19-06-1997

(12) DEMANDE INTERNATIONALE PUBLIÉE EN VERTU DU TRAITÉ DE COOPÉRATION
EN MATIÈRE DE BREVETS (PCT)

(19) Organisation Mondiale de la Propriété
Intellectuelle
Bureau international



(43) Date de la publication internationale
16 août 2001 (16.08.2001)

PCT

(10) Numéro de publication internationale
WO 01/58450 A2

(51) Classification internationale des brevets⁷ :
A61K 31/454, A61P 25/34

(74) Mandataire : THOURET-LEMAITRE, Elisabeth;
Sanofi-Synthelabo, 174, avenue de France, F-75013 Paris
(FR).

(21) Numéro de la demande internationale :
PCT/FR01/00356

(81) États désignés (*national*) : AE, AG, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(22) Date de dépôt international : 7 février 2001 (07.02.2001)

(25) Langue de dépôt : français

(26) Langue de publication : français

(30) Données relatives à la priorité :
00/01682 9 février 2000 (09.02.2000) FR

(84) États désignés (*régional*) : brevet ARIPO (GH, GM, KE,
LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), brevet eurasien
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), brevet européen
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,
MC, NL, PT, SE, TR), brevet OAPI (BF, BJ, CF, CG, CI,
CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Déposant (*pour tous les États désignés sauf US*) :
SANOFI-SYNTHELABO [FR/FR]; 174, avenue de
France, F-75013 Paris (FR).

Publiée :

— sans rapport de recherche internationale, sera republiée
dès réception de ce rapport

(72) Inventeurs; et

(75) Inventeurs/Déposants (*pour US seulement*) : BLAN-
CHARD, Jean, Charles [FR/FR]; 107, avenue de
Paris, F-94160 Saint Mande (FR). MENARD, François
[FR/FR]; 53, rue du Four, F-75006 Paris (FR).

En ce qui concerne les codes à deux lettres et autres abrévia-
tions, se référer aux "Notes explicatives relatives aux codes et
abréviations" figurant au début de chaque numéro ordinaire de
la Gazette du PCT.

(54) Title: USE OF CENTRAL CANNABINOID RECEPTOR ANTAGONIST FOR PREPARING MEDICINES DESIGNED TO
FACILITATE SMOKING CESSATION

(54) Titre : UTILISATION D'UN ANTAGONISTE DES RECEPTEURS AUX CANNABINOÏDES CENTRAUX POUR LA PRE-
PARATION DE MÉDICAMENTS UTILES POUR FACILITER L'ARRÊT DE LA CONSOMMATION DE TABAC

(57) Abstract: The invention concerns the use of N-piperidino-5-(chlorophenyl)-1-(2,4-dichlorophenyl)-4-methylpyrazole-3-car-
boxamide to facilitate smoking cessation.

(57) Abrégé : L'invention concerne l'utilisation du N-piperidino-5-(4-chlorophenyl)-1-(2,4-dichlorophenyl)-4-méthylpyrazole-3-
carboxamide, pour faciliter l'arrêt de la consommation de tabac.




WO 01/58450 A2

UTILISATION D'UN ANTAGONISTE DES RECEPTEURS AUX
CANNABINOÏDES CENTRAUX POUR LA PREPARATION DE MEDICAMENTS
UTILES POUR FACILITER L'ARRET DE LA CONSOMMATION DE TABAC.

La présente invention concerne une nouvelle utilisation d'un antagoniste des récepteurs aux cannabinoïdes centraux dits récepteurs CB1. Plus particulièrement, l'invention se rapporte à l'utilisation d'un antagoniste des récepteurs CB1 pour la préparation de médicaments utiles pour faciliter l'arrêt de la consommation de tabac.

Des familles de composés ayant une affinité pour les récepteurs aux cannabinoïdes ont été décrites dans plusieurs brevets et demandes de brevets, en particulier la demande européenne EP-576 357, qui décrit des dérivés du pyrazole, et la demande WO 96/02248 qui décrit notamment des dérivés du benzofurane.

Plus particulièrement, le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, ci-après dénommé composé A, de formule :



ses sels pharmaceutiquement acceptables et leurs solvats, sont décrits dans le brevet européen EP-656 354 et par M. Rinaldi-Carmona et al. (FEBS Lett., 1994, 350, 240-244), comme antagonistes des récepteurs centraux CB₁.

Il est décrit que le composé A et ses sels qui sont des antagonistes des récepteurs centraux aux cannabinoïdes peuvent être utilisés pour le traitement des troubles de l'appétit, notamment en tant qu'anorexigène, et dans le traitement des troubles liés à l'utilisation de substances psychotropes. De plus, la demande internationale WO99/00119 divulgue l'utilisation des antagonistes des récepteurs aux cannabinoïdes centraux pour traiter les troubles de l'appétence, c'est-à-dire réguler les désirs de consommation, en particulier pour la consommation de sucres, de carbohydrates, d'alcool ou de drogues et plus généralement d'ingrédients appétissants.

On a maintenant trouvé que le composé A, ses sels pharmaceutiquement acceptables et leurs solvats permettent de faciliter l'arrêt de la consommation de tabac,

qu'ils sont utiles dans le traitement de la dépendance à la nicotine et/ou dans le traitement des symptômes du sevrage à la nicotine.

Ainsi, l'administration du composé A, d'un de ses sels pharmaceutiquement acceptables ou solvats, permet d'observer chez des consommateurs de nicotine tels que les fumeurs de tabac, une abstinence tabagique totale ou partielle de façon précoce ou retardée. De plus, les symptômes du sevrage à la nicotine sont très sensiblement atténués voire supprimés, et la prise de poids consécutive à l'arrêt de la consommation tabagique est réduite ou inexistante.

Selon un des ses aspects, la présente invention concerne l'utilisation du N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, d'un de ses sels pharmaceutiquement acceptables ou d'un de leurs solvats pour la préparation de médicaments utiles pour faciliter l'arrêt de la consommation de tabac, dans le traitement de la dépendance à la nicotine et/ou dans le traitement des symptômes du sevrage à la nicotine.

Selon la présente invention, on peut également utiliser le composé A, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats, en association avec un autre principe actif, pour la préparation de médicaments utiles pour faciliter l'arrêt de la consommation de tabac, dans le traitement de la dépendance à la nicotine et/ou dans le traitement des symptômes du sevrage à la nicotine.

Par exemple le composé A peut être associé

- à la nicotine, un agoniste nicotinique ou un agoniste nicotinique partiel, ou bien
- à un inhibiteur de monoamine oxydase (IMAO),
- ou à tout autre principe actif ayant démontré une efficacité pour faciliter l'arrêt de la consommation de tabac, par exemple un antidépresseur tel que le bupropion, la doxépine, la nortriptyline ou un anxiolytique tel que la buspirone, ou bien la clonidine.

Pour son utilisation en tant que médicament, le composé A, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats, seul ou en association avec un autre principe actif, doit être formulé en composition pharmaceutique.

Ainsi la présente invention a également pour objet des compositions pharmaceutiques contenant en association le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats et un autre principe actif, l'autre principe actif étant un composé utile pour faciliter l'arrêt de la consommation de tabac, et/ou utile dans le traitement de la dépendance à la nicotine et/ou dans le traitement des symptômes du sevrage à la nicotine. Ledit autre principe actif étant préférentiellement choisi parmi :

- la nicotine, un agoniste nicotinique ou un agoniste nicotinique partiel, ou bien

- un inhibiteur de monoamine oxidase (IMAO),

- ou tout autre principe actif ayant démontré son efficacité pour faciliter l'arrêt de la consommation de tabac, par exemple un antidépresseur tel que le bupropion, la doxépine, la nortriptyline ou un anxiolytique tel que la buspirone, ou bien la clonidine.

5 Dans les compositions pharmaceutiques de la présente invention pour l'administration orale, sublinguale, sous-cutanée, intramusculaire, intraveineuse, transdermique, locale ou rectale, le principe actif, seul ou en association avec un autre principe actif, peut être administré sous forme unitaire d'administration, en mélange avec des supports pharmaceutiques classiques, aux animaux et aux êtres humains. Les
10 formes unitaires d'administration appropriées comprennent les formes par voie orale telles que les comprimés, les gélules, les pilules, les poudres, les granules, les gommes à mâcher et les solutions ou suspensions orales, les formes d'administration sublinguale et buccale, les aérosols, les implants, les formes d'administration locale, transdermique, sous-cutanée, intramusculaire, intraveineuse, intranasale ou
15 intraoculaire et les formes d'administration rectale.

Dans les compositions pharmaceutiques de la présente invention, le principe actif ou les principes actifs sont généralement formulés en unités de dosage. L'unité de dosage contient 0,5 à 300 mg, avantageusement de 5 à 60 mg, de préférence de 5 à 40 mg par unité de dosage, pour les administrations quotidiennes, une ou plusieurs fois
20 par jour.

Bien que ces dosages soient des exemples de situations moyennes, il peut y avoir des cas particuliers où des dosages plus élevés ou plus faibles sont appropriés, de tels dosages appartiennent également à l'invention. Selon la pratique habituelle, le dosage approprié à chaque patient est déterminé par le médecin selon le mode
25 d'administration, l'âge, le poids et la réponse dudit patient.

Lorsque l'on prépare une composition solide sous forme de comprimés, on peut ajouter au(x) principe(s) actif(s) micronisé(s) ou non un agent mouillant tel que le laurylsulfate de sodium et on mélange le tout avec un véhicule pharmaceutique tel que la silice, l'amidon, le lactose, le stéarate de magnésium, le talc ou analogues. On peut
30 enrober les comprimés de saccharose, de divers polymères ou d'autres matières appropriées ou encore les traiter de telle sorte qu'ils aient une activité prolongée ou retardée et qu'ils libèrent d'une façon continue une quantité prédéterminée de principe actif.

On obtient une préparation en gélules en mélangeant le principe actif ou les principes actifs avec un diluant tel qu'un glycol ou un ester de glycérol et en incorporant le mélange obtenu dans des gélules molles ou dures.
35

Une préparation sous forme de sirop ou d'éllixir peut contenir le principe actif ou les principes actifs conjointement avec un édulcorant, acalorique de préférence, du méthylparaben et du propylparaben comme antiseptiques, ainsi qu'un agent donnant du goût et un colorant approprié.

5 Les poudres ou les granules dispersibles dans l'eau peuvent contenir le principe actif ou les principes actifs en mélange avec des agents de dispersion ou des agents mouillants, ou des agents de mise en suspension, comme la polyvinylpyrrolidone ou polyvidone, de même qu'avec des édulcorants ou des correcteurs du goût.

10 Pour une administration rectale, on recourt à des suppositoires qui sont préparés avec des liants fondant à la température rectale, par exemple du beurre de cacao ou des polyéthylèneglycols.

15 Pour une administration parentérale, on utilise des suspensions aqueuses, des solutions salines isotoniques ou des solutions stériles et injectables qui contiennent des agents de dispersion et/ou des agents solubilisants pharmacologiquement compatibles, par exemple le propylèneglycol ou le butylèneglycol.

20 Ainsi, pour préparer une solution aqueuse injectable par voie intraveineuse on peut utiliser un cosolvant, par exemple un alcool tel que l'éthanol ou un glycol tel que le polyéthylèneglycol ou le propylèneglycol, et un tensioactif hydrophile tel que le polysorbate 80. Pour préparer une solution huileuse injectable par voie intramusculaire, on peut solubiliser le principe actif par un triglycéride ou un ester de glycérol.

Pour l'administration transdermique, on peut utiliser des patches sous forme multilaminée ou à réservoir dans lequel le principe actif est en solution alcoolique.

25 Le principe actif ou les principes actifs peuvent être formulés également sous forme de microcapsules ou microsphères, éventuellement avec un ou plusieurs supports ou additifs.

Le principe actif ou les principes actifs peuvent être également présentés sous forme de complexe avec une cyclodextrine, par exemple α -, β - ou γ - cyclodextrine, 2-hydroxypropyl- β -cyclodextrine ou méthyl- β -cyclodextrine.

30 Parmi les formes à libération prolongée utiles dans le cas de traitements chroniques, on peut utiliser des implants. Ceux-ci peuvent être préparés sous forme de suspension huileuse ou sous forme de suspension de microsphères dans un milieu isotonique.

35 De façon préférentielle, le composé A est administré par la voie orale, en une prise unique par jour.

Selon un autre aspect de l'invention, le composé A, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats et l'autre principe actif associé peuvent être administrés de manière simultanée, séparée ou étalée dans le temps pour faciliter l'arrêt de la consommation de tabac.

5 On entend par "utilisation simultanée" l'administration des composés de la composition selon l'invention compris dans une seule et même forme pharmaceutique.

On entend par "utilisation séparée" l'administration, en même temps, des deux composés de la composition selon l'invention chacun compris dans une forme pharmaceutique distincte.

10 On entend par "utilisation étalée dans le temps" l'administration successive, du premier composé de la composition selon l'invention, compris dans une forme pharmaceutique, puis, du deuxième composé de la composition selon l'invention, compris dans une forme pharmaceutique distincte.

15 Dans le cas de cette "utilisation étalée dans le temps", le laps de temps écoulé entre l'administration du premier composé de la composition selon l'invention et l'administration du deuxième composé de la même composition selon l'invention n'excède généralement pas 24 heures.

20 Les formes pharmaceutiques, comprenant soit un seul des composés constitutifs de la composition selon l'invention soit l'association des deux composés, qui peuvent être mises en œuvre dans les différents types d'utilisations décrites ci-dessus, peuvent par exemple être appropriées à l'administration orale, nasale, parentérale ou transdermique.

25 Aussi, dans le cas d'une "utilisation séparée" et d'une "utilisation étalée dans le temps", deux formes pharmaceutiques distinctes peuvent être destinées à la même voie d'administration ou à une voie d'administration différente (orale et transdermique ou orale et nasale ou parentérale et transdermique etc).

30 L'invention concerne donc également une trousse pour faciliter l'arrêt de la consommation de tabac contenant le composé A et un autre principe actif facilitant l'arrêt de la consommation de tabac dans laquelle ledit composé A et ledit principe actif sont dans des compartiments distincts et dans des conditionnements semblables ou différents, et sont destinés à être administrés de manière simultanée, séparée ou étalée dans le temps. Ledit principe actif est préférentiellement choisi parmi :

- la nicotine, un agoniste nicotinique ou un agoniste nicotinique partiel, ou bien
- un inhibiteur de monoamine oxidase (IMAO),

- ou tout autre principe actif ayant démontré une efficacité pour faciliter l'arrêt de la consommation de tabac, par exemple un antidépresseur tel que le bupropion, la doxepine, la nortriptyline ou un anxiolytique tel que la buspirone, ou bien la clonidine.

5 Selon un autre de ses aspects, l'invention concerne aussi une méthode pour faciliter l'arrêt de la consommation de tabac qui consiste à administrer à un consommateur de nicotine une quantité thérapeutiquement efficace du composé A, d'un de ses sels pharmaceutiquement acceptables ou un de leurs solvats.

10 Les effets du composé A ont été étudiés chez le rat sur un modèle prédictif des effets sur la dépendance envers la nicotine : l'autoadministration de nicotine selon W.T. Corrigan and al. Psychopharmacology, 1989, 99, 473-478.

Le composé A, administré à la dose de 0,3 mg/kg et 1 mg/kg diminue de façon statistiquement significative le nombre d'injections de nicotine, chez des rats qui ont appris à s'autoadministrer de la nicotine par voie intraveineuse.

Ainsi, on a constaté les effets positifs du composé A sur ces 2 modèles.

15 Une étude en double aveugle a été réalisée avec des sujets fumant plus de 15 cigarettes par jour et montrant des symptômes de dépendance à la nicotine. Les patients reçoivent 40 mg du composé A par jour pendant 10 semaines dont 2 semaines avant le début de la période d'abstinence tabagique. On observe dans le groupe traité un taux d'abstinence plus important que dans le groupe recevant un placebo, 20 notamment lors des 4 dernières semaines de traitement. L'abstinence tabagique est confirmée par la mesure hebdomadaire des taux de monoxyde de carbone expiré et de cotinine plasmatique.

EXEMPLE 1 : gélule dosée à 5 mg de composé A.

25	Composé A micronisé	5,00 mg
	Amidon de maïs	51,00 mg
	Lactose monohydrate	99,33 mg
	Polyvidone	4,30 mg
	Laurylsulfate de sodium	0,17 mg
	Carboxyméthyl cellulose de sodium réticulée	8,50 mg
30	Eau purifiée : Q.S. pour granulation humide	
	Stéarate de magnésium	1,70 mg
		<hr/>
	Pour une gélule blanc opaque n° 3 remplie à	170 mg

EXEMPLE 2 : gélule dosée à 10 mg de composé A.

	Composé A micronisé	10,00 mg
	Amidon de maïs	51,00 mg
	Lactose monohydrate	94,33 mg
5	Polyvidone	4,30 mg
	Laurylsulfate de sodium	0,17 mg
	Carboxyméthyl cellulose de sodium réticulée	8,50 mg
	Eau purifiée : Q.S. pour granulation humide	
	Stéarate de magnésium	1,70 mg
10		<hr/>
	Pour une gélule blanc opaque n° 3 remplie à	170 mg

EXEMPLE 3 : gélule dosée à 20 mg de composé A.

	Composé A micronisé	20,00 mg
15	Amidon de maïs	51,00 mg
	Lactose monohydrate	84,33 mg
	Polyvidone	4,30 mg
	Laurylsulfate de sodium	0,17 mg
	Carboxyméthyl cellulose de sodium réticulée	8,50 mg
20	Eau purifiée : Q.S. pour granulation humide	
	Stéarate de magnésium	1,70 mg
		<hr/>
	Pour une gélule blanc opaque remplie à	170 mg

EXEMPLE 4 : comprimé dosé à 10 mg de composé A.

25	Composé A micronisé	10,00 mg
	Amidon de maïs	50,00 mg
	Lactose monohydrate 200 mesh	211,50 mg
	Hydroxypropylméthylcellulose 6 cP	9,00 mg
30	Carboxyméthylamidon sodique	15,00 mg
	Laurylsulfate de sodium	1,50 mg
	Stéarate de magnésium	3,00 mg
	Eau purifiée : Q.S.	
35	Pour un comprimé terminé à	<hr/> 300 mg

EXEMPLE 5 : comprimé dosé à 30 mg de composé A.

	Composé A micronisé	30,00 mg
	Amidon de maïs	80,00 mg
	Lactose monohydrate 200 mesh	252,00 mg
5	Povidone K 30	12,00 mg
	Carboxyméthylcellulose sodique réticulée	20,00 mg
	Laurylsulfate de sodium	2,00 mg
	Stéarate de magnésium	4,00 mg
	Eau purifiée : Q.S.	<hr/>
10	Pour un comprimé terminé à	400 mg

REVENDEICATIONS

1. Utilisation du N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, d'un de ses sels pharmaceutiquement acceptables
5 ou d'un de leurs solvats pour la préparation de médicaments utiles pour faciliter l'arrêt de la consommation de tabac, dans le traitement de la dépendance à la nicotine et/ou dans le traitement des symptômes du sevrage à la nicotine.
2. Utilisation selon la revendication 1 du N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, d'un de ses sels
10 pharmaceutiquement acceptables ou d'un de leurs solvats en association avec un autre principe actif, dans laquelle l'autre principe actif est utile pour faciliter l'arrêt de la consommation de tabac et/ou utile dans le traitement de la dépendance nicotinique et/ou utile dans le traitement du sevrage à la nicotine.
3. Composition pharmaceutique contenant en association le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses
15 sels pharmaceutiquement acceptables ou un de leurs solvats et un autre principe actif dans laquelle l'autre principe actif est utile pour faciliter l'arrêt de la consommation de tabac, et/ou utile dans le traitement de la dépendance à la nicotine et/ou dans le traitement des symptômes de sevrage à la nicotine.
- 20 4. Composition pharmaceutique selon la revendication 3 contenant le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats et la nicotine, un agoniste nicotinique ou un agoniste nicotinique partiel.
- 25 5. Composition pharmaceutique selon la revendication 3 contenant le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats et un inhibiteur de monoamine oxydase.
- 30 6. Trousse pour faciliter l'arrêt de la consommation de tabac contenant le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats et un autre principe actif facilitant l'arrêt de la consommation de tabac dans laquelle le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses sels pharmaceutiquement acceptables
35 ou un de leurs solvats et ledit principe actif sont dans des compartiments distincts et dans des conditionnements semblables ou différents et sont destinés à être administrés de manière simultanée, séparée ou étalée dans le temps.

7. Trousse selon la revendication 6 contenant le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats et la nicotine, un agoniste nicotinique ou un agoniste nicotinique partiel
- 5 8. Trousse selon la revendication 6 contenant le N-pipéridino-5-(4-chlorophényl)-1-(2,4-dichlorophényl)-4-méthylpyrazole-3-carboxamide, un de ses sels pharmaceutiquement acceptables ou un de leurs solvats et un inhibiteur de la monoamine oxydase.

THIS PAGE BLANK (USPTO)